



# **Impact Assessment of Titanium Dioxide Nanoparticles (TiO<sub>2</sub>-NPs) on physico-Chemical Properties of Soil after Harvesting of Cauliflower (*Brassica oleracea* var. *botrytis*) at Prayagraj District of Eastern Uttar Pradesh, India**

**Mohit Bhardwaj<sup>a\*</sup>, Ram Bharose<sup>a</sup>, Tarence Thomas<sup>a</sup>  
and Narendra Swaroop<sup>a</sup>**

<sup>a</sup> *Department of Soil Science and Agricultural Chemistry, Naini Agriculture Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj, Uttar Pradesh 211 007, India.*

## **Authors' contributions**

*This work was carried out in collaboration among all authors. Author MB designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors RB and TT checked the manuscript. Author NS managed the literature cited. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/IJECC/2023/v13i92331

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/102316>

**Original Research Article**

**Received: 25/04/2023**

**Accepted: 29/06/2023**

**Published: 14/07/2023**

\*Corresponding author: E-mail: hdmohit0@gmail.com;

## ABSTRACT

Cauliflower (*Brassica oleracea* var. *botrytis* L.) is one of the most important winter vegetables among the Cole crops which belongs to the genus *Brassica* of the family Brassicaceae. The current research entitled, effect of titanium dioxide nanoparticles (TiO<sub>2</sub>-NPs) on soil physico-chemical characteristics in cauliflower (*Brassica oleracea* var. *botrytis*) at Prayagraj District of Eastern Uttar Pradesh was conducted during 2022-2023 on the Crop Research Farm of the Department of Soil Science and Agricultural Chemistry, Prayagraj (U.P). There were seven different treatments combinations included in the study comprising treatment T<sub>1</sub> as control and treatment T<sub>2</sub> to T<sub>7</sub> contains 100% RDF through chemical fertilizers along with TiO<sub>2</sub> application as seed treatment and foliar spray with concentration of 500, 1000 and 1500 ppm that were replicated three times in Randomized Block Design. The results demonstrated that available N, P & K were significantly affected and soil physical parameters (Bulk density, particle density and water holding capacity) were non-significantly affected by application of TiO<sub>2</sub> nanoparticles applied as seed treatment or foliar spray.

**Keywords:** Cauliflower; titanium dioxide; seed treatment; foliar spray; nanoparticles.

## 1. INTRODUCTION

The soil is a natural habitat for plant growth and it gives plants nutrients. Some soils are productive and sustain luxuriant plant growth with very little human effort, whereas others may be unproductive and support absolutely no beneficial plant life despite all human effort. For soil to be productive, it must meet the following criteria: (i) be easily tillable and fertile; (ii) have all necessary nutrients in quantities that are readily available to plants; and (iii) be physically sound enough to sustain plants and have the ideal moisture and air content for healthy root development. Throughout the plant's existence, the soil must continuously provide these necessities.

Vegetables are essential for human nourishment. These foods are an essential group because they offer the body with vitamins, minerals, protein, carbs, and fiber in addition to their nutritional worth and therapeutic properties. One of the most significant winter vegetables among the many vegetables is cauliflower (*Brassica oleracea* var. *botrytis* L.), which is a member of the genus *Brassica* and family Brassicaceae. Cauliflower may be grown in a variety of soil types, including clay and loam, but the most ideal conditions are reasonably deep loam soil, adequate fertility, and a decent growing environment [1-4]. Because plants in light soil are more susceptible to drought, water stress has a negative impact on the formation of curds. Soils having a high moisture-holding capacity are preferable for late season/summer. Light soils are necessary for early crops, whereas loamy and clay loam soils are better suited

for midseason and late maturing varieties [5-9].

As cauliflower is mainly a crop that can withstand low temperatures, it does best in cool, humid climates [10-12]. The fifth-largest vegetable crop in India is cauliflower, which is largely cultivated in the winter. India produces 5873.3 thousand MT/ha of cauliflower over an area of 433.9 thousand ha, with a productivity rate of 19.8 MT/ha. Fertilizers have a critical function, as seen by the sharp rise in vegetable output and the rise in fertilizer use. West Bengal produced the most cauliflower in 2014, with 73.6 thousand hectares and 1879.0 thousand MT/ha, respectively (National Horticulture Board, 2014).

The experimental side of nanotechnology is gradually giving way to applications in real world settings. The development of so-called "smart fertiliser" as new facilities to improve nutrient use efficiency and lower costs of environmental protection has been made feasible by nanotechnology [13-15]. These materials can be used as fertilizer transporters or controlled-release vectors. Active food ingredients at the nanoscale (1–100 nm) are the focus of nano fertilizers. These engineered nanoparticles (ENPs) can enter the cells and leaves of plants and can also carry chemicals and DNA into the cells of plants.

In comparison to commonly used pesticides and their formulations, the advantages of nanomaterial-based formulations include improved efficacy due to higher surface area to volume ratio, which provides a better opportunity for interaction, higher solubility, induction of

systemic activity due to smaller particle size and higher mobility, and lower toxicity due to the elimination of organic solvents. Nanotechnology creates novel materials that have the potential to overcome many of the drawbacks of traditional conventional materials. It is also predicted to soon become a major economic force with a great potential for attaining sustainable agriculture, particularly in underdeveloped nations. In the field of agriculture, the use of nanomaterial is comparatively new and needs further exploration. Moreover, the emphasis is given to the recent developments in plant science that focuses on agricultural practices, plant growth and yield.

The reason TiO<sub>2</sub> NPs were chosen is that they are among the most frequently discussed types in the literature that discusses employing NPs in agriculture. Certain applications include nano fertilizers and nano pesticides to track goods and nutrient levels to boost productivity without decontaminating soils and waters and protection against pests and diseases are of major interest to the agricultural industry. The ingredient Titanium-dioxide (TiO<sub>2</sub>), one of the most often used in the agricultural and energy sectors, is thought to be advantageous for plant growth and development.

There is some study that primarily focuses on the benefits of nanoparticles for agriculture. The improvement of some key element concentrations in plant tissues, an increase in yield, and an increase in the chlorophyll content of paprika are three of Titanium's most significant effects on plants (*Capsicum annuum* L.). Nano TiO<sub>2</sub> application boosted soybean germination, growth, and nitrate reductase activity. Additionally, it has been observed that TiO<sub>2</sub> nanoparticles enhance spinach (*Spinacia oleracea*) seed germination and plant growth of spinach. The application of nanoparticles as

fertilizer holds considerable promise for improving crop growth and reducing environmental risks. Manufactured nanoparticles may be able to store nutrients on their surfaces for use by organisms as a source of nutrients.

Even, when applied as foliar spray also improved crop production of Cluster bean (*Cyamopsis tetragonoloba* L.). Plants may quickly absorb nano-formulated fertilizers, and they may have a prolonged effective duration of nutrient delivery in the soil or on the plant.

## 2. MATERIALS AND METHODS

The experiment was carried out at Crop Research Farm, Department of Soil Science and Agricultural Chemistry, Naini Agriculture Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj (UP). The different amounts of Titanium-dioxide (TiO<sub>2</sub>) were given as a foliar spray and a seed treatment (500, 1000, 1500 ppm). There were three replications and 7 treatment combinations, with treatment T<sub>1</sub> serving as the control. Treatments T<sub>2</sub>-T<sub>7</sub> comprised of 100% RDF (chemical fertilizers) along with seed treatment and foliar spray of 500, 1000 and 1500 ppm TiO<sub>2</sub> nanoparticles for cauliflower.

## 3. RESULTS AND DISCUSSION

The data given in Table 2, clearly revealed that bulk density of soil is influenced by various treatments. The application of inorganic and organic source of nutrients along with TiO<sub>2</sub> nanoparticles had non-significant effect on bulk density of soil. The range of values of bulk density varies from 1.14 to 1.22 Mg m<sup>-3</sup>. Among various treatments the maximum bulk density (1.22 Mg m<sup>-3</sup>) was recorded in treatment T<sub>4</sub> and T<sub>3</sub> followed by treatment T<sub>2</sub> and T<sub>4</sub>. The minimum bulk density (1.14 Mg m<sup>-3</sup>) was

**Table 1. Methods for the determination of different physical properties of soil**

Particular	Method employed	Reference(s)
Bulk density	Graduated measuring cylinder	Muthuaval et al., [16].
Particle density	method.	
Water holding capacity		
Available Nitrogen (kg/ha)	Alkaline potassium permanganate method	Subbiah and Asija, 1956
Available Phosphorus (kg/ha)	Calorimetric method	Olsen et al., 1954
Available Potassium (kg/ha)	Flame photometric method	Toth and Prince, 1949

noted under treatment T<sub>5</sub>. A scrutiny of data presented in Table 2 revealed that application of nanoparticles with chemical fertilizers had non-significant effect on particle density of soil. The range of particle density varies from 2.43 to 2.62 Mg m<sup>-3</sup>. However, the maximum particle density (2.62 Mg m<sup>-3</sup>) was recorded under treatment T<sub>2</sub> comprising 100% RDF (chemical fertilizers) + 500 ppm TiO<sub>2</sub> (foliar spray) in cauliflower and the minimum particle density (2.43 Mg m<sup>-3</sup>) was noted under treatment T<sub>6</sub> [100% RDF (chemical fertilizers) + 1000 ppm TiO<sub>2</sub> (seed treatment)] and treatment T<sub>7</sub> (100% RDF + 1500 ppm TiO<sub>2</sub> seed treatment). The data regarding the water holding capacity in soil as influenced by different treatments is given in Table 2, the application of various treatments had non-significant effect on water holding capacity of soil in cauliflower. The range of water holding capacity varies from 46.20 to 46.81 %. However, the maximum water holding capacity (46.81%) was recorded under treatment T<sub>3</sub> i.e., 100% RDF (chemical fertilizers) +1000 ppm TiO<sub>2</sub> as foliar spray. The minimum (46.20 %) water holding capacity was noted under treatment T<sub>6</sub> with 100% RDF (chemical fertilizers) + 1000 ppm TiO<sub>2</sub> (seed treatment). Similar findings are in accordance with the studies conducted by Badhulkar et al. [17], Selvi et al. [18] and Bajpai et al. [19], concluded that long term research and experimentation may modify some of the physicochemical characteristics of soil.

A scrutiny of data presented in Table 3 revealed that there was significant increment in available soil nitrogen due to application of nanoparticles along with chemical fertilizers. The significant higher (287.74 kg ha<sup>-1</sup>) value of available nitrogen was recorded under treatment T<sub>4</sub> comprising 100% RDF (chemical fertilizers) +

1500 ppm TiO<sub>2</sub> (foliar spray) in cauliflower followed by treatment T<sub>1</sub>. However, the minimum available nitrogen (239.49 kg ha<sup>-1</sup>) was noted under treatment T<sub>6</sub> [100% RDF (chemical fertilizers) + 1000 ppm TiO<sub>2</sub> (seed treatment)].

The data regarding the available phosphorus in soil as influenced by different treatments is given in Table 3. On statistical analysis it was noted that significantly higher (22.60 kg ha<sup>-1</sup>) phosphorus content of soil was revealed under treatment T<sub>4</sub> i.e., 100% RDF (chemical fertilizers) +1500 ppm TiO<sub>2</sub> as foliar spray in cauliflower. The minimum (16.06 kg ha<sup>-1</sup>) amount of available phosphorus was noted under treatment T<sub>6</sub> with 100% RDF (chemical fertilizers) + 1000 ppm TiO<sub>2</sub> (seed treatment) as given in Table 3.

A significant difference was revealed on analysing the data regarding available potassium content in soil as given in Table 3. Significantly higher (202.11 kg ha<sup>-1</sup>) value of available potassium was recorded in treatment T<sub>4</sub> in cauliflower. The minimum (192.95 kg ha<sup>-1</sup>) available potassium was recorded under treatment T<sub>6</sub> with 100% RDF (chemical fertilizers) + 1000 ppm TiO<sub>2</sub> (seed treatment) in cauliflower.

Nitrogen, Phosphorus and Potassium availability in treatments with foliar application and seed treatment with TiO<sub>2</sub> nanoparticles may have been further reduced due to improved nutrient absorption from the soil brought on by application of different treatments. Similar research was also reported by Osterhies (1998), who came to the conclusion that nutrient supplementation through foliar and seed treatment might improve soil nutrient absorption, boost crop productivity, and decrease the quantity of fertilizer used on the soil (Ahmad, 1988).

**Table 2. Effect on bulk density, particle density and water holding capacity of soil in cauliflower by the application of nanoparticles (TiO<sub>2</sub>)**

Treatments	Bulk density (Mg m <sup>-3</sup> )	Particle density (Mg m <sup>-3</sup> )	Water holding capacity (%)
T1 Control (100% RDF)	1.19	2.55	46.49
T2 100% RDF + 500 ppm TiO <sub>2</sub> Foliar Spray	1.20	2.62	46.67
T3 100% RDF + 1000 ppm TiO <sub>2</sub> Foliar Spray	1.22	2.56	46.81
T4 100% RDF + 1500 ppm TiO <sub>2</sub> Foliar Spray	1.22	2.51	46.26
T5 100% RDF + 500 ppm TiO <sub>2</sub> Seed Treatment	1.14	2.58	46.53
T6 100% RDF + 1000 ppm TiO <sub>2</sub> Seed Treatment	1.16	2.43	46.20
T7 100% RDF + 1500 ppm TiO <sub>2</sub> Seed Treatment	1.14	2.55	46.80
F-Test	NS	NS	NS
S. Em. (±)	-	-	-
C.D. at 5%	-	-	-

**Table 3. Effect of TiO<sub>2</sub> (nano-particles) and chemical fertilizers on available Nitrogen, phosphorus and potassium (kg ha<sup>-1</sup>) of soil after crop harvest**

Treatments	Available N (Kg ha <sup>-1</sup> )	Available P (Kg ha <sup>-1</sup> )	Available K (Kg ha <sup>-1</sup> )
T1 Control (100% RDF)	278.69	21.60	199.37
T2 100% RDF + 500 ppm TiO <sub>2</sub> Foliar Spray	265.25	20.29	196.87
T3 100% RDF + 1000 ppm TiO <sub>2</sub> Foliar Spray	258.84	18.48	197.67
T4 100% RDF + 1500 ppm TiO <sub>2</sub> Foliar Spray	287.74	22.60	202.11
T5 100% RDF + 500 ppm TiO <sub>2</sub> Seed Treatment	241.94	17.01	196.59
T6 100% RDF + 1000 ppm TiO <sub>2</sub> Seed Treatment	239.49	16.06	192.95
T7 100% RDF + 1500 ppm TiO <sub>2</sub> Seed Treatment	247.64	17.68	198.63
<b>F-Test</b>	S	S	S
<b>S. Em. (±)</b>	0.35	0.26	0.51
<b>C.D. at 5%</b>	1.06	0.8	1.58

#### 4. CONCLUSIONS

The study revealed that seed treatment with nanoparticles (TiO<sub>2</sub>) and foliar spray in combination with chemical fertilizers had a significant effect on available NPK (kg ha<sup>-1</sup>) of soil after crop harvest and a non-significant effect on bulk density (Mg m<sup>-3</sup>), particle density (Mg m<sup>-3</sup>) and water holding capacity of soil. TiO<sub>2</sub> nanoparticle treatments with foliar spraying and seed treatment resulted in increased nutrient uptake from the soil, which further decreased soil-available macronutrient availability. This could explain why there was a decrease in the primary macronutrients that were available in the soil in these treatments.

#### ACKNOWLEDGEMENT

For providing the essential research facilities, the authors are grateful to the Department of Soil Science and Agricultural Chemistry, Naini Agriculture Institute, Sam Higginbottom University of Agriculture Technology and Sciences Prayagraj, Uttar Pradesh.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Chinnamuthu CR, Boopathi PM. Nanotechnology and Agroecosystem. Madras Agricultural Journal. 2009;96:17-31.
- Galbraith DW. Nanobiotechnology: Silica breaks through in plants. Nature Nanotechnology. 2007;2(5):272-273.
- Naderi MR, Danesh-Shahraki A. Nanofertilizers and their roles in sustainable agriculture. International Journal of Agricultural and Crop Sciences. 2013;5(19):2229-2235.
- De Rosa MC, Monreal C, Schnitzer M, Walsh R, Sultan Y. Nanotechnology in fertilizers. Nature Nanotechnology. 2010; 5(2):91.
- Fraceto LF, Grillo R, De Medeiros GA, Scognamiglio V, Rea G, Bartolucci C. Nanotechnology in agriculture: Which innovation potential does it have? Frontiers in Environmental Science. 2016;4:20.
- Hruby M, Cigler P, Kuzel S. Titanium in plant nutrition: The contribution to understanding the mechanism of titanium action in plants. Journal of Plant Nutrition. 2002;25:577-598.
- Zheng L, Hong F, Lu S, Liu C. Effect of nano-TiO<sub>2</sub> on strength of naturally aged seeds and growth of spinach. Biological Trace Element Research. 2005;105:83-91.
- Vakhrouchev AV, Golubchikov VB. Numerical investigation of the dynamics of nanoparticle systems in biological processes of plant nutrition. Journal of Physics: Conference Series 2007;61(1): 31-35.
- Navarro E, Baun A, Behra R, Hartmann NB, Filser J, Miao A, Quigg A, Santschi PH, Sigg L. Environmental behavior and ecotoxicity of engineered nanoparticles to algae, plants, and fungi. Ecotoxicology. 2008;17:372-386.
- Gogos A, Knauer K, Bucheli TD. Nanomaterials in plant protection and fertilization: Current state, foreseen applications, and research priorities. Journal of Agricultural and Food Chemistry. 2012;60(39):9781-9792.

11. Anupam M, Sinha I, Das R. Application of Nanotechnology in Agriculture: Future Prospect, paper presented in conference at Mumbai; 2015.
12. Chaudhary I, Singh V. Titanium Dioxide Nanoparticles and its Impact on Growth, Biomass and Yield of Agricultural Crops under Environmental Stress: A Review. *Research Journal of Nanoscience and Nanotechnology*. 2020;10:1-8.
13. Raliya R, Tarafdar JC. ZnO Nanoparticle Biosynthesis and Its Effect on Phosphorous-Mobilizing Enzyme Secretion and Gum Contents in Clusterbean (*Cyamopsis tetragonoloba* L.). *Agricultural Research* 2013;2:48–57.
14. Rameshaiah GN, Jpallavi S. Nano fertilizers and nano sensors—an attempt for developing smart agriculture. *International Journal of Engineering Research and General Science*. 2015;3(1):314-320.
15. Black CA. *Methods of Soil Analysis*. Part I and II. American Society of Agronomy, Inc., Madison, Wisconsin, U. S. A. 1965;1-2:1572.
16. Muthuaval PC, Udaysooriyan R, Natesa PP. Ramaswami. Introduction to Soil Analysis, Tamil Nadu Agriculture University, Coimbatore-641002.
17. Badhulkar RM, Wandile WP, Balpande SS. Residual effect of long-term application of FYM and fertilizer on soil properties (Vertisols) and yield of soybean. *Journal of the Indian Society of Soil Science*. 2000;48:89-92.
18. Selvi D, Santhy P, Dhakshinamoorthy M, Maheshwari M. Microbial population and biomass in rhizosphere as influenced by continuous intensive cultivation and fertilization in an inceptisol. *Journal of the Indian Society of Soil Science*. 2004;52: 254-257.
19. Bajpai RK, Shrikant C, Upadhyay SK, Urkurkar JS. Long term studies on soil physico-chemical properties and productivity of rice-wheat system as influenced by integrated nutrient management in inceptisol of Chattisgarh. *Journal of the Indian Society of Soil Science*. 2006;54:24-29.

© 2023 Bhardwaj et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/102316>