



Effect of Nutrient Management on Growth, Yield and Quality of Onion (*Allium cepa* L.)

Hema Solanki ^{a*}, Vishwanath ^a, Jyoti Diwakar ^a and Manvendra Singh ^a

^a *Department of Horticulture, Raja Balwant Singh College, Bichpuri, Agra (U.P) 283105, India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i193608

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104651>

Original Research Article

Received: 13/06/2023

Accepted: 17/08/2023

Published: 26/08/2023

ABSTRACT

A field experiment was conducted on the Horticulture Research Farm of the RBS College, Bichpuri, Agra during the *rabi* season of 2016-18. In this experiment eight treatments were tested with three replications in randomized block design (RBD). Treatments were (T1) recommended dose of fertilizer (RDF) (T2) 100% RDF + Azotobacter, (T3) 100% RDF + FYM, (T4) 100% RDF + Azotobacter + FYM, (T5) 80% RDF + Azotobacter + FYM, (T6) 80% RDF + FYM, (T7) 80% RDF + Azotobacter and (T8) control. The application of 100%RDF +Azotobacter + FYM (T4) significantly maximum plant height (50.14cm), maximum number of green leaves (8.65), maximum leaf length(45.53cm). maximum fresh weight of top of plant 45.34gm), maximum fresh weight of bulbs (84.32gm), maximum diameter of bulb (6.9cm) and the highest bulb yield of 33.95 t/ha. were recorded from treatment T4 (100%RDF +Azotobacter + FYM)

Keywords: *Nutrient management; azotobacter; Farm Yard Manure (FYM); yield and quality; onion (*Allium cepa* L.).*

*Corresponding author: E-mail: solankihema1995@gmail.com;

1. INTRODUCTION

Onion (*Allium cepa* L.) is the most important commercial vegetable and spice crop cultivated extensively in India. The primary center of origin is Central Asia [1]. It belongs to Amaryllidaceae family having $2n = 16$ chromosome number, biennial herb type plant with tabular leaves and bulbs are formed by the attachment of swollen leaf bottom to underground part of stem, which is small and rudimentary. Its flower type is 'umbel,' which have many small flowers. This is cross pollinated crop. Onion has many medicinal properties like anti-flatulence and cure dysentery, reducing cholesterol levels, it may help in reducing high blood pressure and also useful to prevent hair fall. India is second largest producer of the world. The randomly use of chemicals results in bad soil health, erosion loss of organic matter, nitrate pollution and also health hazard from human beings. The bio-fertilizers are alternative source to meet the requirement of crops and to bridge the further gaps and also get the sustainable production and productivity as well as quality. Azotobacter is an inorganic fertilizer and it is free living, aerobic nitrogen fixing and non-symbiotic bacteria. It fixes about 20 kg/ha atmospheric nitrogen in the roots of the many crops like vegetables, spices and condiments besides the pulses and enhance the germination (Sethi and Adhikary 2012). FYM is basically made by decomposition of animal dung, urine other dairy waste, dry leaves and straw waste. Farm yard manure have all macro and micro nutrients having 0.5 % N, 0.2% P₂O₅ 0.5%-K₂O. To reducing uses of chemical fertilizer in the soil, to save soil health and promote crop quality, longevity of soil, FYM is very affective. FYM less expensive than chemical fertilizers. Just few researchers namely Banjare et. al. [2] and Talwar Dilpreet et. al. [3] studied in this regard to find out the effect of nutrient management with bio-fertilizers on the onion crop. However, till now no systematic approaches are available so far made to utilize the agro-ecological condition of Gangatic plains of Uttar Pradesh and small information is available about the organic cultivation of this crop in the country. Therefore, it was considered worthwhile to carry out the present investigation on the growth yield and quality of onion cv.– Agree found light red (AGFR) under Gangatic alluvial soil condition of Uttar Pradesh.

2. MATERIALS AND METHODS

The present investigation was under taken during the *rabi* season of year 2016-17 and 2017-18 for

study the response of different combinations of bio-fertilizer, chemical and FYM (Azotobacter + NPK + FYM) by the vegetative growth, and quality character of onion (*Allium cepa* L.). Experiment was conducted at the Agricultural Research Farm of Raja Balwant Singh college Bichpuri, Agra (U.P) The soil of the experimental field was Gangatic with calcareous layer at the depth of about 1.0 – 1.5 meters, it was slightly alkaline in reaction pH 7.84 and well drained. Weather conditions of Agra was semi-arid sub-tropical climate with hot dry summer and hardy cold winters. Temperature falls at about 1-2-°C in winter and increased about maximum at 45-47°C. in summer. The treatments were (T1) Recommended dose of fertilizer (RDF) (T2) 100% RDF + Azotobacter (T3) 100% RDF + FYM, (T4) 100% RDF + Azotobacter + FYM, (T5) 80% RDF + Azotobacter + FYM, (T6) 80% RDF + FYM, (T7) 80% RDF + Azotobacter and (T8) control. The eight treatments were replicated three times in randomized block design in 1.60m x 1.20m size plots. The recommended dose of nitrogen, phosphorus and potash were applied at the time of transplanting. The half dose of nitrogen was applied as basal and in two parts after 30 and 60-days after transplanting (DAT). farm yard manure was applied 15 days before transplanting and the bio-fertilizer (Azotobacter) was applied as seedling dipping treatment in Azotobacter and water solution at 2kg/ha. All the required cultural operations along with the irrigation was done as per requirement of the crop. The all growth studies recorded at 15 and 30 DAT. The bulb was harvested at the mature stage. The experimental site is gangatic alluvial in origin with calcareous layer at the depth of about 1.5m -2m, it was slightly alkaline in reaction (pH 7.84). The soil pf experimental field is deficient in total nitrogen, low in organic carbon, medium in phosphorus and rich in potash.

3. RESULTS AND DISCUSSION

The pooled data of length of plant presented in table and figure shows that maximum length of plant was 23.59, 39.98, 50.14 and 23.34cm at 30, 60, 90 DAT, respectively and at harvest were found to be significantly higher in T4 (100% RDF + Azotobacter + FYM), which was significantly followed by T3 (100% RDF + FYM) and T5 (80 % RDF + Azotobacter + FYM), the lower height of plant was observed in T8 (control). These finding are in agreement with the findings of Jawadagi et. al. [4] and Kurrey et. al. (2018).




Fig. 1. Effect of nutrient management on the vegetative growth of onion




Fig. 2. Effect of nutrient management on the vegetative growth of onion

Table 1. Effect of nutrient management on the growth attributed of onion

Treatment	Plant height (cm)	No. of leaves /plant	Length of leaf	Width of leaf (cm)	Diameter of pseudo-stem (cm)	Weight. of tops (gm)
T ₁	43.53	6.55	38.23	1.51	1.43	41.73
T ₂	44.31	6.66	39.66	1.65	1.30	41.85
T ₃	47.85	7.66	42.46	1.88	1.53	43.65
T ₄	50.14	8.65	45.53	2.00	1.76	45.34
T ₅	46.25	7.55	41.10	1.82	1.33	42.56
T ₆	44.53	7.22	40.36	1.73	1.30	42.18
T ₇	41.3	6.11	37.25	1.44	1.30	41.34
T ₈	39.52	5.88	33.96	1.28	1.20	38.31
S.Em±	0.656	0.193	0.330	0.051	0.708	0.714
CD(P=0.05)	5.97	0.588	1.00	0.155	0.214	2.16

Table 2. Effect of nutrient management on the yield and quality attributes of onion

Treatment	Bulb diameter (cm)	Weight of bulb (gm)	Bulb yield qt/ha	Specific gravity
T ₁	4.05	68.32	222.67	1.08
T ₂	4.21	71.57	250.93	1.04
T ₃	4.93	82.05	301.48	1.05
T ₄	5.21	84.32	339.58	1.62
T ₅	4.96	80.13	282.43	1.10
T ₆	4.47	77.13	268.64	1.04
T ₇	4.60	66.68	218.72	0.89
T ₈	4.04	63.40	193.13	0.919
S.Em±	0.246	1.27	30.85	0.105
CD(=0.50)	0.747	3.88	93.59	0.319

Number of green leaves at 30, 60, 90, DAT and at harvest were found to be significantly maximum in T₄ (100% RDF + Azotobacter + FYM) that was closely followed by T₃ (100% RDF + FYM) and the lowest number of green leaves per plant was observed in T₈. The possible reason for increased number of green leaves per plant may be due to the improvement in growth related attributes because of certain promoting substances secreted by bio-fertilizers and better uptake of water.

Leaf length is one of the important characters for determining crop vigour and yield. In the present study all the inorganic + bio-fertilizer + organic manure combination treatments were significantly superior to other. The maximum length of longest leaf at 30, 60, 90, DAT and at harvest were found to be significantly higher in T₄ (100% RDF + Azotobacter + FYM) which, was followed by T₃. (100% RDF + FYM) and T₅ (80% RDF + Azotobacter + FYM) whereas, the minimum leaf length was recorded in control (T₈). Similar results were observed by Meena et al. [5] and Tinna et al. (2020)

The width of longest leaf at 30, 60, 90, DAT and harvest were found to be significantly higher in (T₄) 100% recommended dose of fertilizer with the organic manure/ substance and bio-fertilize. Increases in width of leaf with optimum level of chemical fertilizer with organic manures could be due to good growth of plant that enhanced by the production of bio active substance having similar effects as that the growth regulator besides nitrogen fixation through bacterial fertilizers. Similar result observed by Singh et al. [6]

Considerable variation in fresh weight of the above parts of plant with T₄ (100% RDF + Azotobacter + FYM) might be attributed due to the improvement in overall growth and development of plants as led to higher (45.34gm) fresh weight of top of plant and lower (38.31gm) fresh weight was in control (T₈). This enabled the plant to draw more nutrients from soil for proper growth and development as stated by Talwar et al. [7].

Significantly higher bulb diameter was noted under T₄ (100% RDF + Azotobacter + FYM) and

lower was found in control (T8). Similar results observed by Brinjh et al. [8].

Average bulb weight was also recorded maximum in the bulbs receiving 100% RDF + Azotobacter + FYM (T4) followed by 100% RDF + FYM (T3). The increases in the bulb weight could be due to the increased uptake of nutrients and buildup of sufficient photosynthesis enabling the increases in size of bulbs ultimately resulting in the increased averaged bulb weight. These results are in confirmation with the finding of Yogita et al. [9] and Kumar et al. (2019).

The maximum specific gravity of bulb (1.626) was in the treatment T4 (100% RDF + Azotobacter + FYM) followed by T5 (80% RDF + Azotobacter + FYM) (1.107) and minimum (0.89) specific gravity was also observed in T7 (100% RDF + Azotobacter). Similar results noted by Bhagwat et al. [10]

The main and important objective of any production is to have maximum crop yield for better returns. The data regarding bulb yield reveals a significant difference among the different treatments. The maximum bulb yield/plot and hectare (5.17 kg and 339.58q) respectively, was obtained in T4 (100% RDF + Azotobacter + FYM) that was statistically at par with T3 and T5 and minimum was noticed in T8. The increase in yield may be due to more number of bulbs per plot, bulb size and average weight of bulb. Use of Azotobacter and FYM, not only makes the atmospheric N available to plants but also enhances the plant growth and bulb yield to release hormones, vitamins and nutrient. Similar findings were also reported by Talwar et al. [7] and Afify et al. [11], [12-15].

4. COCLUISON

All the inorganic, organic and bio-fertilizers combination of treatments were found statically better on performance of onion, on the basis of experiment conducted, it is concluded that the treatment T4 (100% RDF + Azotobacter + FYM) was found superior among all other treatments for growth, yield and quality characters of onion. Further trial of this research work in different location of the Uttar Pradesh are needed to recommended results at farmer's level.

CONFERENCE DISCLAIMER

Some part of this manuscript was previously presented in the conference: 6th International

Conference on Strategies and Challenges in Agricultural and Life Science for Food Security and Sustainable Environment (SCALFE-2023) on April 28-30, 2023 in Himachal Pradesh University, Summer Hill, Shimla, HP, India. Web Link of the proceeding: <https://www.shobhituniversity.ac.in/pdf/Souvenir-Abstract%20Book-Shimla-HPU-SCALFE-2023.pdf>

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Vavilov NI. Origin, variation, immunity and breeding of cultivated plants. Journal of Chronical Botany. 1951;13(1/6):364.
2. Banjare C, Shukla N, Sharma PK, Patanwar M, Chandravanshi D. Effect of organic substances on yield and quality of onion (*Allium cepa* L.). International Journal of Farm Sciences. 2015;5(1):30-35.
3. Dilpreet Talwar. () Influence of biofertilizers on growth, yield, quality and nutrient uptake in onion (*Allium cepa* L.) Indian Journal of Ecology. 2016;43(1):124-27.
4. Jawadagi RS, Basavaraj N, Patil BN, Hemlta Naik B. Effect of different sources of nutrients on growth, yield and quality of onion (*Allium cepa* L) Cv. Bellary red. Karnataka Journal of Agricultural Science. 2012;25(2):232-35.
5. Meena AK, Paliwal R, Meena KK. Effect of organic manures and bio-fertilizers on growth and quality attributes of kharif onion (*Allium cepa* L.) in semi-arid region. Indian Research Journal of Genetics and Biotechnology. 2015;7(1):73-76.
6. Singh A, Ram RB, Meena ML. Efficacy of different sources of nutrients and bio-fertilizers on growth yield and quality of onion. International Research Journal of Natural and Applied Sciences. 2015;2(10):64-70.
7. Talwar P, Gupta NK, Dubey S. Effect of organic, inorganic and biofertilizers on the growth and productivity of garlic (*Allium sativum* L) cv. G-323. Crop Research. 2012;43(1,2 & 3):89-97.
8. Brinjh S, Kumar S, Kumar D, Kumar M. Effect of integrated nutrient management on the growth, yield and quality in onion cv.

- Pusa Madhavi. Plant Archives. 2014;14(1): 557-59.
9. Yogita, Ram RB. Effect of chemical and bio-fertilizers on quality of onion. (*Allium cipa* L). Hort Flora Research Spectrum. 2012;1(4):367-70.
 10. Bhagwat KP, Shinde KG, Bhalekar MN, Yadav DB. Effect of organic manure, inorganic fertilizers and biofertilizers on growth and yield attributes in onion (*Allium cepa* L). 2016;13(1):13-15.
 11. Aida H Afify, FIA. Hauka and El-Sawah, A.M. (2018) Plant Growth-Promoting Rhizobacteria enhance onion (*Allium cepa* L) Productivity and minimize requiste chemical fertilization. Env. Biodiv. Soil Security. 2: 119-29
 12. Amit K, Meena ML, Shirvran BC, Harvindar Pal, Meena BL. Impact of bio-fertilizer on growth, yield and quality of onion (*Allium cepa* L) cv. Pusa Red. Plant Archives. 2019;19(1):772-76
 13. Devendra KK, Maheendra KL, Gaurav SP. Effect of azotobcter on growth and yield of onion (*Allium cepa* L.) Journal of Pharmacognosy and Phytochemistry. 2018;7(1):1171-75.
 14. Deeksha T, Naveen G, Sandeep S, Gulab P, Neena C. Utilization pf Plant growth promoting rhizobacteria as root dipping of seedlings for improving bulb yield and curtailing mineral fertilizer use in onion under field conditions. Scientia Horticulturae. 2020;(270):109432.
 15. Santosh KK, Siba PA. Azotobacter: A plant growth-promoting rhizobacteria used as biofertilizer. Global Science Book. 2012:68-74.

© 2023 Solanki et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/104651>