



Importance of Bio-Priming on Improving Seed Quality in Chilli (*Capsicum Annuum* L.)

Shivayogi Ryavalad ^{a++}, Ashok Surwenshi ^{b#*},
Arvind Kumar Rathod ^{c++}, Muske Deepa N. ^{d++},
Srinivas N. ^e and Rajkumar M. ^{f#}

^a RHREC, Dharwad, Karnataka, India.

^b CPH, COH, BIDAR, Karnataka, India.

^c SST, COH, BIDAR, Karnataka, India.

^d GPB, COH, BIDAR, Karnataka, India.

^e FSC, COH, BIDAR, Karnataka, India.

^f HEEU, YADGIR, Karnataka, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i224160

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/109787>

Original Research Article

Received: 20/09/2023

Accepted: 24/11/2023

Published: 29/11/2023

ABSTRACT

India grows a lot of chilli (*Capsicum annuum* L.), and one of its most valuable commodities is its seed. However, the quality of the seeds is lost if they are stored for the months between harvest and the subsequent sowing. The seeds of Rudra and Byadgi Dabbi varieties were subjected to bio-priming with the *Pseudomonas fluorescens*, *Bacillus* and *Trichoderma viride* on qualitative parameters with seven treatments. The experiment was carried out using CRBD. The culture of

⁺⁺ Assistant Professor;

[#] Associate Professor;

*Corresponding author: E-mail: surwenshi@gmail.com;

Pseudomonas fluorescens, *Bacillus* and *Trichoderma viride* at 2.5g/l and 5.0g/l concentration were used as pre-sowing bio-priming treatments. From the results it was found that the Bio-priming influenced significantly on the capability of the chilli seedlings with the improved germination percentage of 98.69% (2.5 g *P. fluorescens*, T₁), shoot length 6.53cm (5 g *P. fluorescens*, T₂), root length 7.18cm (5 g *T. viride*, T₆), seedling length 13.43 cm (5 g *P. fluorescens*, T₂) and seedling vigour index 1216.44 (5 g *P. fluorescens*, T₂) in Byadgi Dabbi cultivar and where as in Rudra cultivar recorded germination percentage of 99.75% (T₂), shoot length 4.13cm (5 g *T. viride*, T₆), root length 7.38cm (5 g *T. viride*, T₆), seedling length 11.56 cm (5 g *T. viride*, T₆) and seedling vigour index 1140.12 (5 g *T. viride*, T₆) when compared with control. Therefore, bio-priming with *Pseudomonas fluorescens*, *Bacillus* and *Trichoderma viride* are effective in enhancing the quality parameters of seed in chilli.

Keywords: Bio-priming; *Pseudomonas fluorescens*; *Trichoderma viride*; *Bacillus*; chilli.

1. INTRODUCTION

Among solanaceous vegetables, the chilli (*Capsicum annum* L.) is a major crop grown in subtropical and tropical regions. Ripe and unripe fruits are utilized for a variety of industrial and therapeutic purposes [1]. The main causes of the decreased production and inferior quality characteristics of chilli include imbalanced fertilizer usage, inadequate plant protection, poor development, and sub-optimal plant population [2]. Poor and inconsistent germination usually leads to a suboptimal plant population. Invigorating seeds to increase germination rate, increase development uniformity, and shorten the time it takes for many vegetables and certain field crops to emerge has been the subject of several research in recent years. Nowadays, seed priming is a popular commercial technique that increases seedling uniformity and speeds up germination in a variety of crops [3]. The most important ingredient for agriculture is seed. Therefore, it is essential to preserve the quality of the seeds using naturally occurring treatments that offer several advantages. One of the most significant advancements in pre-sowing procedures is seed priming, which promotes uniform and quick growth and growth of seeds while also enhancing their resistance to unfavorable climatic circumstances [4]. According to Galhaut et al. [5], bio-priming is the most straightforward, cost-effective, and environmentally friendly method of delivering beneficial microorganisms to the agroecosystem among seed priming strategies. Therefore, understanding how bio-priming can be applied to improve plant nutrition, restore soil health, and maintain crop/seed quality is crucial to achieving the UN Sustainable Development Goals, which include ending hunger, poverty, and promoting good health and well-being [6]. In order to preserve seed, a procedure known as "bio-

priming" combines the physiological part of disease control—seed hydration—with the natural component of disease control—seed inoculation with helpful organisms. It is an ecological strategy that combats soil- and seed-borne infections by employing either bacteria or specific fungal antagonists. Certain bacteria and fungi, most notably *Pseudomonas fluorescens*, *Bacillus*, and *Trichoderma viride*, are known to inhibit illness and promote plant development.

A fungus called *Trichoderma viride* is employed as a bioagent. It is applied to seeds and soil to control a range of illnesses brought on by fungi. Many crops, including beans (*Phaseolus vulgaris*), cucumbers (*Cucumis sativus*), peppers (*Capsicum annum*), carnations (*Dianthus carophyllus*), maize (*Zea mays*), and wheat (*Triticumaestivum*), have been shown to have an enhanced growth response when exposed to *Trichoderma* sp [7]. *Trichoderma* create phytohormones that are linked to or produce cytokinin-like molecules, such as zeatyn and gibberellin GA3. These phytohormones, whose metabolic routes have been discovered [8], include indol acetic acid (IAA) and ethylene (M. Ahsanur Rahman et al. 2012). Improved bio-fertilization may result from the regulated synthesis of these substances [8]. Another common bioagent for disease control is *pseudomonas fluorescens*, also referred to as plant growth rizobacteria because of its ability to produce phytohormones, phyto-growth-regulating substances (PGRs), mineralize organic phosphorus, produce phytoalexin-like compounds, enhance mineral absorption, and more [9]. When compared to the control, *Pseudomonas* sp. MML2212 and *Pseudomonas fluorescens* were found to increase dry weight and plant height on rice and green gram [10,11,12]. In order to assess how several

genotypes of chillies responded to bio-priming with *Pseudomonas fluorescens* and *Trichoderma viride* in terms of growth, yield, and quality features, the current study was started.

2. MATERIALS AND METHODS

The present study was carried out at UHS, Bgalkote using byadgi dabbi and rudra genotypes of chilli. Which are very popular in the area. The research was carried out using complete randomized design comparing with seven treatments like T₁- 2.5 g *P. fluorescens*, T₂-5 g *P. fluorescens*, T₃-2.5 g *Bacillus*, T₄-5 g *Bacillus*, T₅- 2.5 g *T. viride*, T₆- 5 g *T. viride*., T₇- Untreated control.

2.1 Seed Treatment

Seeds of both Byadgi Dabbi and Rudra genotypes of chilli were bio-primed with *P. fluorescens*, *Bacillus* and *T. viride* individually for twenty-four hours at different concentration and As a control, the seeds were steeped in distilled water for the same amount of time. The seeds were used for lab tests after being shade-dried for 30 minutes following the treatment.

2.2 Laboratory Experiments

For the in vitro experiment, two layers of Whatman filter paper soaked in sterilized distilled water were put on petriplates lined with 100 seeds, with four replications for each genotype and treatment, and incubated at 25±10C. According to ISTA, the germination percentage was measured at (14 Days After Sowing-DAS). After 14 DAS, five seedlings at random from each replication were chosen to measure the length of the roots and shoots using the Paper Plate technique. The information was logged and subjected to a rigorous analysis.

3. RESULTS AND DISCUSSION

Following pre-sowing bio-priming, the seeds were subjected to laboratory testing to evaluate their quality using several criteria such as vigor index, root length, and shoot length, seedling length, and germination %. The results revealed that there was significant effect of different treatment on the quality parameters' of chilli.

3.1 Germination Percentage

The result revealed that the highest germination percentage was observed in 2.5 g *P. fluorescens* (T₁) with value of 98.69% for after 14 DAS of sowing followed by 5 g *Bacillus* (T₄), 5 g *T. viride* (T₆), 2.5 g and *T. viride* (T₅) respectively whereas lowest was recorded in treatment 2.5 g *Bacillus* (T₃) of about 90.00% for germination percentage in Byadgi Dabbi variety. The Rudra variety recorded higher germination percentage in 5 g *P. fluorescens* (T₂) with value of 99.75% and all other treatments were on par with each other with lower germination percentage recorded in untreated control (T₇) of about 98.00% mentioned in Tables 1 and 2. It was discovered that *Pseudomonas fluorescens* had a higher average effect on this characteristic in both types compared to *Bacillus* and *Trichoderma viride*. The Rudra variety responded similarly to the other varieties when pre-sowing bio-priming of seed germination, and the impact of the three bio-inoculants was nearly as good as that of the control. In contrast, substantial variations were seen in Byadgi Dabbi across all treatments. In all conditions, *Trichoderma viride* also enhanced the germination %; data to corroborate this were reported by Hanson [13], Mishra and Sinha [14], and Mukhtar I [15].

3.2 Root Length

Root length was recorded as maximum as 7.18cm and 6.93 cm in 5 g *T. viride* (T₆) and (T₂) in Byadgi Dabbi variety, when average was made over treatments, though 5 g *T. viride* (T₆) and 5 g *P. fluorescens* (T₂) carried out comparably statistically for this character. The untreated control group (T₇) had the significantly shortest root length, with 2.5 g *Bacillus* (T₃) and 5 g *Bacillus* (T₄) following closely behind. The distinct genetic expression might be one of the factors. The genotype Rudra recorded as maximum as 7.38 cm and 7.25 cm in 5 g *T. viride* (T₆) and 2.5 g *P. fluorescens* (T₁) treatments whereas shortest root length was recorded for Untreated control (T₇) of about 4.80cm followed by 2.5 g *Bacillus* (T₃) with root length of about 4.90cm (Table. 1 and Table 2). *Pseudomonas fluorescens*, *Bacillus*, and *Trichoderma viride* bioinoculant all showed a significant beneficial effect over control when it came to the development of this trait in both kinds. Comparable outcomes with reference to the root growth of cucumber, loofah, and bitter gourd from screened *Trichoderma* strains were discovered by Lo, C.T., and Lin, CY. (2002).

Table 1. Effect of seed bio-priming on seed quality of Byadgi Dabbi chilli genotypes

Treatments	Germination Percent (%)	Root Length (cm)	Shoot Length (cm)	Seedling Length (cm)	Vigour Index
T ₁	98.69	5.38	4.33	9.71	957.99
T ₂	90.75	6.93	6.53	13.43	1216.44
T ₃	90.00	4.63	4.75	9.38	845.02
T ₄	97.25	4.88	5.15	10.03	974.78
T ₅	96.13	6.05	4.21	10.26	986.58
T ₆	96.25	7.18	2.30	9.48	911.99
T ₇	94.50	4.53	4.70	8.23	777.26
SEM	1.15	0.09	0.11	0.21	20.89
CD@5%	3.37	0.27	0.32	0.61	61.44
CV	2.42	3.26	4.88	4.13	4.38

Table 2. Effect of seed bio-priming on seed quality of Rudra chilli genotypes

Treatments	Germination Percent (%)	Root Length (cm)	Shoot Length (cm)	Seedling Length (cm)	Vigour Index
T ₁	99.50	7.25	3.03	10.28	1022.37
T ₂	99.75	5.90	2.15	8.05	803.09
T ₃	99.38	4.90	2.53	7.43	738.01
T ₄	99.44	6.03	3.23	9.25	919.94
T ₅	98.50	5.20	3.43	8.63	849.54
T ₆	99.15	7.38	4.13	11.50	1140.12
T ₇	98.00	4.80	3.88	8.68	850.00
SEM	0.32	0.13	0.07	0.22	22.48
CD@5%	0.93	0.38	0.21	0.66	66.12
CV	0.64	4.34	4.54	4.90	4.98

3.3 Shoot Length

The longest shoot was produced in Byadgi Dabbi variety with treatment 5 g *P. fluorescens* (T₂) of about 6.53cm when overall performance of the all treatments were considered followed by 5 g *Bacillus* (T₄) with 5.15cm whereas shortest shoot was recorded in 5 g *T. viride* (T₆) with 2.30cm. In Rudra cultivar longest shoot length noticed with 4.13cm in (T₆) and shortest recorded of about 2.15cm in 5 g *P. fluorescens* (T₂). It was discovered that the average effects of *Bacillus*, *Trichoderma viride*, and *Pseudomonas fluorescens* were positive and statistically significant when compared to the unprimed control. While *Pseudomonas fluorescens* was seen in the Byadgi Dabbi variety, a larger effect of *Trichoderma viride* was detected in the Rudra variety. The Rudra cultivar's untreated control (T₇) reacted to priming similarly to *Bacillus*, *Pseudomonas fluorescens*, and *Trichoderma viride*.

3.4 Seedling Length

Significantly longest seedlings were produced by 5 g *P. fluorescens* (T₂) with 13.43cm followed by

(T₂) with 10.26cm in Byadgi Dabbi, while shortest recorded for untreated control (T₇) with 8.23cm. In Rudra variety longest seedlings were produced by 5 g *T. viride* (T₆) with 11.50cm followed by 2.5 g *P. fluorescens* (T₁) with 10.28cm whereas shortest recorded for 2.5 g *T. viride* (T₅) with 8.63cm. *Bacillus* performed statistically similarly in both genotypes. Regardless of the cultivars, the untreated control (T₇) generated seedlings of the most tiny kind. Over time, it was shown that the average untreated control of *Bacillus*, *Trichoderma viride*, and *Pseudomonas fluorescens* was significantly positive compared to the unprimed control. However, in the Rudra variety, *Trichoderma viride* was observed to have a bigger impact at higher concentrations. A table showing the change in control due to priming may also be used to evaluate the similar trend that was observed for each specific variety and was more pronounced following the bio-priming of seeds with *Trichoderma viride* Rai et al. [1], Sarkar et al. [6].

3.5 Vigour Index

Similar to seedling length, the untreated control group (T₇) exhibited the lowest magnitude of

vigour index. This might be attributed to both the increased effect of its seed germination potential and the inherited potential of its seedling length. In both types, bio-priming had a significant impact on the increase in vigor index compared to control (Rai et al. [1], Sarkar et al. [6], Asaduzzaman et al. [16], Mishra, D.S., and Sinha, A.P. 2000). In Rudra cultivar, *Trichoderma viride* (5 g T. viride, T₆=1140.12) had a greater impact than *Pseudomonas fluorescens* (2.5 g P. fluorescens, T₁=1022.37 and 5 g P. fluorescens, T₂=803.09) for determining the average vigor index. whereas its reverse trend positive superiority in influence of *Pseudomonas fluorescens* (5 g P. fluorescens, T₂=1216.44) could be noticed over that of *Trichoderma viride* (5 g T. viride, T₆=911.99) in Byadgi dabbi variety (Table 1 and Table 2), Although both types' responses to bio-priming for the development of seeds with increased vigor above control were statistically equal, they followed a similar pattern. It was found that all of the bio-inoculants had very identical effects on both types [17,18]. exception was noted for 2.5 g *Bacillus* (T₃) in for which low superiority in influence with low concentration of *Trichoderma viride*.

4. CONCLUSION

From the study it was found that there is enhancement in the seed quality parameters of chilli when there is bioprimgng of *Pseudomonas fluorescens*, *Bacillud* and *Trichoderma viride*. The significant variation was found when compared with the control treatment, so from the study it concluded that to improve the crop quality there is need to have the bioprimgng of the mentioned microorganisms. Preferably *Pseudomonas fluorescens* to be used for the better results when compared with others.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Rai AK, Behera S. Response of seed bio-priming on chilli (*Capsicum annum* L.) Journal of Pharmacognosy and Phytochemistry. 2019;8(1):900-903.
- Alam MZ, Hamim I, Ali MA, Ashrafuzzaman M. Effect of Seed Treatment on Seedling Health of Chilli. J Environ. Sci. & Natural Resources. 2014; 7(1):177-181
- Basra SMA, Zia MN, Mehmood T, Afzal I, Khaliq A. Comparision of different invigoration techniques in wheat (*Triticum aestivum* L.) seeds. Pak J Arid Agr. 2003;5:11-17.
- Dhanya BA. Evaluation of microbial seed priming in relation to seed germination, plant growth promotion in *Morinda citrifolia* L. (Noni). M. Sc. (Ag.) Thesis. University of Agricultural Sciences, Bangalore; 2014.
- Galhaut L, Lespinay A, Walker DJ, Bernal MP, Correal E, Lutts S. Seed priming of *Trifolium repens* L. improved germination and early seedling growth on heavy metal-contaminated soil. Water Air Soil Pollution. 2014;225:1-15.
- Sarkar D, Singh S, Parihar M, Rakshit A. Seed bio-priming with microbial inoculants: A tailored approach towards improved crop performance, nutritional security, and agricultural sustainability for smallholder farmers. Current research on environmental sustainability. 2021;3: 100093.
- Lo CT, Lin CY. Screening strains of *Trichoderma* spp. for plant growth enhancement in Taiwan. Plant pathology Bull. 2002;11:215-220.
- Osiewacz HD. Genes, mitochondria, and aging in filamentous fungi. Ageing Res. Rev. 2002;28.
- Mukerji KG, Manoharachary C, Singh J. Microbial activity in the rhizosphere. Springer, New York. Verlag, Berlin. 2006;155-172.
- Mathivanan N, Prabhavathy VR, Vijayanandraj VR. Application of Talc Formulations of *Pseudomonas fluorescens* Migula and *Trichoderma viride* Pers. ex S.F. Gray Decrease the Sheath Blight Disease and Enhance the Plant Growth and Yield in Rice. J Phytopathol. 2005; 153:697-701.
- Shanmugaiah V. Biocontrol potential of Phenazine -1- carboxamide producing plant growth promoting rhizobacterium *Pseudomonas aeruginosa* MML2212 against sheath blight disease of rice. Ph.D. Thesis, University of Madras, Chennai, India; 2007.
- Shanmugaiah V, Ramesh S, Jayaprakashvel M, Mathivanan N. Biocontrol and plant growth promoting potential of *Pseudomonas* sp. MML2212 from the rice rhizosphere. In: Proceedings of the 1st Int. Symposium on B.C.B.P.D., stadt, Germany; 2005.

13. Hanson LD. Reduction of Verticillium wilt symptoms in cotton following seed treatment with Trichoderma virens. J Cotton. Sci. 2000;4:224-231.
14. Mishra DS, Sinha AP. Plant growth promoting activity of some fungal and bacteria agents on rice seed germination and seedling growth. Tropical Agric. 2000;77:188-191.
15. Mukhtar I. Influence of Trichoderma species on seed germination in okra. Mycopath. 2008; 6(1&2):47-50
16. Asaduzzaman M, Alam MJ, Islam MM, Effect of Trichoderma on seed germination and seedling parameters of chilli. Journal of the J Sci. Foundation. 2010;8(1&2):141-150.
17. Dubey SC, Suresha M, Singha B. Evaluation of Trichoderma species against Fusarium oxysporum f. sp. ciceris for integrated management of chickpea wilt. Biological Control. 2007; 40:118-127.
18. Harris D, Joshi A, Khan PA, Gothkar P, Sodhi PS. Onfarm seed priming in semiarid agriculture: development and evaluation in maize, rice and chickpea in India using participatory methods. Exp. Agric.1999;35:15-29.

© 2023 Ryavalad et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/109787>