



Clinico-Radio-Histopathological Aspects of a Compound Odontome with Impacted Deciduous Canine: An Uncommon Case Report

Anwasha Paul ^{a++}, Sudeshna Bagchi ^{a#*},
Sk. Abdul Mahmud ^{a†}, Swapan Kumar Purkait ^{b‡}
and Rudra Prasad Chatterjee ^{a†}

^a Department of Oral and Maxillofacial Pathology, Guru Nanak Institute of Dental Sciences and Research, Kolkata, India.

^b Department of Oral and Maxillofacial Pathology, Buddha Institute of Dental Sciences & Hospital, Patna, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jammr/2024/v36i105598>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/123885>

Case Report

Received: 15/07/2024

Accepted: 17/09/2024

Published: 21/09/2024

⁺⁺ MDS PGT;

[#] Senior Lecturer;

[†] Professor;

[‡] Professor & Head;

*Corresponding author: E-mail: sudeshna.bagchi2@gmail.com, sudeshna.bagchi@gnidsr.ac.in;

Cite as: Paul, Anwasha, Sudeshna Bagchi, Sk. Abdul Mahmud, Swapan Kumar Purkait, and Rudra Prasad Chatterjee. 2024. "Clinico-Radio-Histopathological Aspects of a Compound Odontome With Impacted Deciduous Canine: An Uncommon Case Report". *Journal of Advances in Medicine and Medical Research* 36 (10):142-48. <https://doi.org/10.9734/jammr/2024/v36i105598>.

ABSTRACT

Odontomes are hamartomatous lesions of odontogenic origin, classified under 'Benign Mixed Epithelial & Mesenchymal Odontogenic Tumors' by WHO 2022. They are mostly detected as incidental findings on standard radiographs due to their slow growth and benign behaviour. Occasionally, they may cause complications such as delayed tooth eruption, impaction, deflection, or retention of teeth. Only 2% cases have shown impaction of a deciduous tooth. We hereby report a case of a compound odontoma in a 5-year-old child causing impaction of a deciduous canine.

Keywords: Compound Odontome; Primary dentition; Maxilla; Impacted deciduous.

1. INTRODUCTION

The term "Odontome" was first coined by French physician Pierre Paul Broca in 1866 in his monograph 'Traite Des Tumeurs' to designate any tumour arising from odontogenic tissues [1]. However, with increasing cognizance, the usage of the term has gradually shifted and Odontomes are now considered as hamartomatous lesions of odontogenic origin, classified under 'Benign Mixed Epithelial & Mesenchymal Odontogenic Tumors' by WHO 2022 [2]; accounting for about 20% to 67% of all odontogenic neoplasms [3,4].

Odontomas can occur at any age, but are typically identified within the first two decades of life [5]; with the mean age of detection being 14.8 years [6]. There is no gender predilection [3]. Normally, odontomas are small, but on rare occasions, they can enlarge and cause expansion of bones [5]. While the majority of odontomas are asymptomatic and are incidental findings on standard radiographs, a small number can even result in complications such as delayed tooth eruption, impaction, deflection, retention of deciduous teeth or infrabony defects [7]. Odontomes are common in the permanent dentition but rarely documented in conjunction with primary teeth [8,9]. The exact etiology of odontoma is not quite apparent [6] and has been associated with a number of local or genetic factors [3].

The WHO categorizes odontomes into compound and complex types [10]. Compound odontomes are typically seen in the incisor cuspid region of the maxilla, whereas complex odontomes are more abundant in the molar and premolar regions of the mandible [11]. Compound odontomas demonstrate tooth-like features on radiography, encircled by a narrow radiolucent zone. Histopathologically, they consist of tiny "denticles" with observable morphologic differentiation and nearly normal dentin, cementum, and enamel matrix relationships. In contrast, complex odontomes appear as an irregular calcified mass surrounded

by a thin radiolucent rim. Histopathologically, they are composed of a haphazard array of less differentiated odontogenic tissues, forming an amorphous mass [4].

This report presents a rare case of compound odontoma associated with the primary dentition of a 5-year-old boy.

2. CASE REPORT

2.1 Clinical Findings

A 5-year-old male patient reported to the Department of Oral and Maxillofacial Pathology with the chief complaint of unerupted maxillary left deciduous canine and a gradually enlarging painless hard swelling in the region since the last 3 months. There was no history of trauma to the mouth or face and no significant medical history. Overall, the patient had good general health.

On extraoral examination, no obvious facial asymmetry was detected (Fig. 1 [A]). Intraoral examination revealed the presence of unerupted left maxillary deciduous canine (63) and a fairly delineated, dome-shaped swelling in the left maxillary anterior region extending from the distal surface of left maxillary deciduous lateral incisor (62) to the mesial surface of left maxillary deciduous first molar (64), causing obliteration of labial vestibular sulcus. The overlying mucosa appeared smooth and stretched with normal physiological pigmentation (Fig. 1 [B]). On palpation, the swelling was bony hard in consistency and no tenderness was elicited.

2.2 Diagnostic Assessment

Panoramic radiograph revealed the presence of a well circumscribed, round to ovoid, radiopaque mass measuring about 2 cm x 1.5 cm, encased by a narrow radiolucent zone, in the maxillary left anterior region, obscuring the pathway of eruption of the unerupted primary canine (63). The radiopaque mass resembled a conglomeration of multiple small tooth-like structures (Fig. 2 [A]).



Fig. 1. [A] Extraoral photograph; [B] Intraorally, presence of dome-shaped swelling in the left maxillary anterior region




Fig. 2. [A] OPG showing radiopaque mass resembling multiple small tooth-like structures in the maxillary left anterior region; Reconstructed CT images in Frontal [B1] and Occlusal [B2] 3D view and Panoramic view [B3]; [C1-C4] Reformatted multiplanar images through left maxilla

Cone Beam Computed Tomography (CBCT) of maxilla revealed deciduous dentition in relation to 51, 52, 61, 62, 64, 65; crown formation in relation to 11, 12, 21, 22, 23, 24, 25, 26, 27 with

incomplete root formation; vertical impaction in relation to 63; and an ill-defined, round to oval radiopacity with a well-corticated periphery measuring 23.7 mm (supero-inferiorly) x 17.6

mm (mesio-distally) x 15.5 mm (labio-palatally) noted at the level of crestal bone; extending from the distal aspect of 62 to the mesial aspect of 64, mesio-distally. The internal structure of the pathology was mixed radiopaque-radiolucent in nature, with the multiple small radiopacities resembling teeth. It was surrounded by a radiolucent area and an outer thin radiopaque rim (Fig. 3 [B1-C2]). Expansion and thinning of the labial cortex with minute areas of breach were noted but the palatal cortex was intact. The highest position on the incisal tip of the crown of vertically impacted 63 having a single completely formed root was noted to be approximately 17.1 mm below the alveolar crest and abutting the labial cortex (Fig. 2 [C3 & C4]). The palatal surface of the tooth was noted to be in approximation with left antrum and mesial surface abutted the lateral wall of left nasal cavity. Based on Lekholm & Zarb classification (1985), the bone type was noted to be Type III bone showing thin layer of cortical bone surrounding a core of trabecular bone of good strength.

Based on the clinical and radiographic findings, a provisional diagnosis of Compound Odontoma was made.

2.3 Surgical intervention & Histopathological Evaluation

Surgical excision of the mass in toto was performed under general anaesthesia. Gross specimen resembled a conglomerated mass of malformed tooth-like structures (Fig. 3 [A]).

The specimen was decalcified and processed using formalin fixation and paraffin embedding. Haematoxylin & Eosin-stained sections revealed the presence of tooth-like entities (Fig. 3 [B]). They were composed of relatively homogenous dentinoid material showing dentinal tubule-like structures. Scant enameloid matrix was focally noted. They enclosed clefts containing pulp-like substance and was surrounded by a layer of fibrous connective tissue (Fig. 3 [C-E]).

The overall histopathological features confirmed the diagnosis of "Compound Odontoma".




Fig. 3 [A] Gross specimen; [B] 4x view showing tooth-like entity; [C] 10x view showing dentinoid material enclosing pulp-like tissue; [D] 40x view showing dentinal tubule like structures; [E] 40x view showing enameloid and dentinoid matrix in transverse section




Fig. 4. [A] Post-operative photograph; [B] Post-operative OPG

2.4 Follow Up & Outcome

Patient was advised periodic post-operative follow-up. Proper healing without any serious complications was noted. Patient was referred to Department of Pedodontia for further management and restoration of normal occlusal dynamics (Fig. 4 [A & B]).

3. DISCUSSION

Odontomas are mixed odontogenic tumors, that arise when fully differentiated epithelial and mesenchymal cells proliferate and develop into ameloblasts and odontoblasts. They are mostly composed of enamel and dentine, though variable amounts of pulp and cementum may be present [6,12,13] and are considered as hamartomatous malformations rather than true neoplasms [14].

Various studies have attributed a number of causes to the etiopathogenesis of odontomas, including local trauma, infections and/or inflammatory conditions, remnants of cell rests of Serres, mature ameloblasts, odontoblastic hyperactivity, aberration in genetic control of odontogenesis or manifestations of hereditary anomalies like Gardner's syndrome, Hermann's syndrome among others [3,4,11,15-18]. Hitchin hypothesized that odontomas can arise from a mutation in a gene, a hereditary condition, or interference with the genetic regulation of tooth development, potentially occurring during the postnatal stage [19]. On the other hand, Levy has documented the traumatic injury-induced experimental induction of this lesion in rats [20].

Xavier et al. demonstrated that a non-degradable variant of β -catenin expressed solely in SOX2-positive postnatal dental epithelial stem cells can activate WNT signalling aberrantly enough to

develop malformations resembling odontomas. The mutant β -catenin in oral epithelium also induces ectopic expression of developmental signals like Shh and Bmp4 that impact adjacent tissues and promote abnormal mesenchymal proliferation and condensation. This significantly contributes to the formation of odontoblasts, dentine and pulp consistent with the tooth-like structures of odontomas [21].

Odontomas can be diagnosed at any age, but are most frequently encountered in the first two decades of life [5,7,22]. According to studies by various authors, occurrence of odontomas have a male-to-female ratio ranging from 0.44:1 to 1.5:1 [4]; however, it is generally agreed that no significant gender bias exists [3-5,16].

Compound odontomes are commonly located in the maxilla with a marked propensity for occurrence in the incisor-cuspid region [5-7,23]. They usually tend to remain asymptomatic, but may be associated with delayed eruption or impaction of teeth, and at later stages may occasionally effectuate enlargement of cortical bones and swelling [3]. Similar findings were reported in the case under discussion. Infection or regional lymphadenopathy may also be observed in severe cases.

Odontomes are rarely associated with the deciduous dentition [8,9,15]. In his review of 396 cases, Katz reported only 2% to be associated with the inability of a primary tooth to erupt [24]. The striking feature of the present case was its association with the primary dentition and the consequent impaction of a primary canine.

On radiographic analysis, compound odontomes reveal an irregular radiopaque structure with variable contour and size, composed of multiple tiny, malformed, primitive tooth-like entities,

having a peripheral narrow radiolucent halo [4,14,22].

Histopathological evaluation of compound odontomes show multiple structures having resemblance to miniature, single-rooted teeth comprised of a central pulpal core enclosed within a shell of dentin and partly covered by varying amounts of demineralized enamel matrix and cementum, contained within a capsule of loose fibrous connective tissue [4,14,22]. Identical radiographic and histopathological features were noted in the present case.

Since odontomes are benign encapsulated lesions, complete surgical enucleation is the preferred treatment. There is very little proclivity for recurrence [11,22]. Furthermore, it is crucial to conduct a thorough clinical and radiographic examination of the patient at regular follow-ups to evaluate any irregularity in the subsequent dentition.

4. CONCLUSION

Delay in eruption or exfoliation of primary teeth, or impaction of succedaneous permanent teeth should be meticulously evaluated clinically and radiographically to detect the presence of odontomes. Despite having limited growth potential, odontomes may lead to complications like cystic change or interference with normal occlusion which necessitates its surgical removal and subsequent periodic monitoring. Thus, regular assessment of dentition is of paramount importance as early diagnosis of odontomas permit simpler, less expensive treatment plans and ensures better prognosis.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

CONSENT

As per international standards, parental written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Philipsen HP, Reichart PA. Classification of odontogenic tumours. A historical review. *J Oral Pathol Med.* 2006;35(9):525-9.
2. Soluk-Tekkesin M, Wright JM. The World Health Organization Classification of Odontogenic Lesions: A Summary of the Changes of the 2022 (5th) Edition. *Turk Patoloji Derg* 2022;38(2):168-184.
3. Čabov T, Fuchs PN, Zulijani A, Čabov Ercegović L, Marelić S. Odontomas: Pediatric Case Report and Review of The Literature. *Acta Clin Croat* 2021;60(1):146-152.
4. Boffano P, Zavattero E, Rocchia F, Gallesio C. Complex and compound odontomas. *J Craniofac Surg* 2012;23(3):685-8.
5. Nelson BL, Thompson LD. Compound odontoma. *Head Neck Pathol* 2010;4(4):290-1.
6. Sivapathasundaram B, Preethi S, Rajendran R. Odontogenic Tumors. In: Sivapathasundaram B editors. *Shafer's Textbook of Oral Pathology*. 8th ed. India: Elsevier; 2016;117-8.
7. Isola G, Cicciù M, Fiorillo L, Matarese G. Association between odontoma and impacted teeth. *J Craniofac Surg* 2017;28(3):755-758.
8. Bordini J Jr, Contar CM, Sarot JR, Fernandes A, Machado MA. Multiple compound odontomas in the jaw: case report and analysis of the literature. *J Oral Maxillofac Surg* 2008;66(12):2617-20.
9. Kharbanda OP, Saimbi CS, Kharbanda R. Odontome--a case report. *J Indian Dent Assoc* 1986;58(7):269-71.
10. WHO Classification of Tumours Editorial Board. *Head and neck tumours* [Internet; beta version ahead of print]. Lyon (France): International Agency for Research on Cancer. (WHO classification of tumours series, 5th ed. 2022; 9. Available: <https://tumourclassification.iarc.who.int/chapters/52>.
11. Jain N, Pillai A, Varshini A, Kawadkar D, Dubey K, Singh A. Odontome- Revisted. *J Clinical Research and Reports* 2021;9(2).
12. Machado Cde V, Knop LA, da Rocha MC, Telles PD. Impacted permanent incisors

- associated with compound odontoma. *BMJ Case Rep* 2015 Jan 12;2015:bcr2014208201.
13. Das UM, Viswanath D, Azher U. A compound composite odontoma associated with unerupted permanent incisor: a case report. *Int J Clin Pediatr Dent* 2009;2(2):50-5.
 14. Neville BW, Damm DD, Allen CM, Chi AC. *Odontogenic Cysts and Tumors*. In: *Oral and Maxillofacial Pathology*. 4th ed. Missouri: Elsevier. 2016;674-5.
 15. Satish V, Prabhadevi MC, Sharma R. Odontome: A Brief Overview. *Int J Clin Pediatr Dent*. 2011;4(3):177-85.
 16. Rana V, Srivastava N, Kaushik N, Sharma V, Panthri P, Niranjana MM. Compound Odontome: A Case Report. *Int J Clin Pediatr Dent*. 2019;12(1):64-67.
 17. Sajesh S, Edsors E, Nandagopan S, Devi SS, Praveen V, Chidambareswaran S. Compound and Complex Odontomes: Case Series with Surgical Management and Brief Review of Literature. *J Pharm Bioallied Sci* 2021;13 Suppl 2:S1772-S1777.
 18. Girish G, Bavle RM, Singh MK, Prasad SN. Compound composite odontoma. *J Oral Maxillofac Pathol* 2016;20:162.
 19. Hitchin AD. The aetiology of the calcified composite odontomes. *Br Dent J*. 1971;130(11):475-482.
 20. Levy BA. Effects of experimental trauma on developing first molar teeth in rats. *J Dent Res* 1968;47(2):323-7.
 21. Xavier G, Patist A, Healy C, Pagrut A, Carreno G, Sharpe PT et al. Activated WNT signaling in postnatal SOX2-positive dental stem cells can drive odontoma formation. *Sci Rep* 2015;5:14479
 22. de Oliveira BH, Campos V, Marçal S. Compound odontoma--diagnosis and treatment: three case reports. *Pediatr Dent* 2001;23(2):151-7.
 23. Soluk Tekkesin M, Pehlivan S, Olgac V, Aksakalli N, Alatlı C. Clinical and histopathological investigation of odontomas: review of the literature and presentation of 160 cases. *J Oral Maxillofac Surg*. 2012;70(6):1358-61.
 24. Katz RW. An analysis of compound and complex odontomas. *ASDC J Dent Child* 1989;56(6):445-9.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/123885>