



Oxidative Changes among Women Suffering From Urinary Tract Infection in Enugu Metropolis, Nigeria

Christian Ihechukwu Uzoukwu ^a, Johnpaul Chukwu ^a,
Kingsley Chimuanya Umezurike ^{b*},
Mary-Theodora Barbara Ochiabuto ^b,
Chidera Godson Chukwuemeka ^b,
Blessing Uzoma Nwoye ^b, Edmund Osi Ugomore ^c
and George Okechukwu Chukwuma ^b

^a Department of Medical Laboratory Science, University of Nigeria, Nsukka, Nigeria.
^b Department of Medical Laboratory Science, Nnamdi Azikiwe University, Awka, Nigeria.
^c Department of Microbiology, University of Port Harcourt, Port Harcourt, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMPS/2024/v26i2667

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/102977>

Original Research Article

Received: 15/05/2023
Accepted: 21/07/2023
Published: 17/01/2024

ABSTRACT

Background: The action of bacterial infection in the urinary tract can trigger the release of free radicals which could progress and overwhelm endogenous antioxidants, hence oxidative changes occur. This work aimed to estimate the level of oxidized low density lipoprotein (Ox-LDL) and superoxide dismutase (SOD), among women suffering from urinary tract infection in Enugu metropolis.

*Corresponding author: E-mail: chimunyaaprofkings@gmail.com;

Materials and Methods: Ethical approval was obtained from the Health Research and Ethics committee (HREC) of the University of Nigeria Teaching Hospital (UNTH), Ituku–Ozalla, Enugu State. Informed consent was obtained from all recruited subjects. About forty (40) women diagnosed of UTIs at the UNTH were recruited for the study and twenty (20) women free of UTIs and who are not registered with UNTH were used as a control for this study. Blood specimens were collected and analysed using the ELISA technique to check for Ox-LDL and SOD levels.

Results: The mean value of the antioxidant enzyme, SOD, was lower than that of the control which suggests a possibility of oxidative stress. However, Ox-LDL mean value was lower than that of the control population showing absence of lipid peroxidation in all sampled population, which may have been pronounced if the subjects had complicated UTI like pyelonephritis.

Conclusion: This study showed a drop in the levels of an antioxidant enzyme among UTI patients placing the patients at risk of oxidative stress. Ox-LDL should be higher in test than in control population indicating lipid peroxidation. However, in this study, the Ox-LDL showed no increase in values. Oxidative stress should be looked out for and treated by physicians when treating UTIs.

Keywords: *Urinary tract infection; superoxide dismutase; oxidized low density lipoprotein; oxidative stress.*

1. BACKGROUND

Urinary tract infection is a common contagion among men and women but the incidence is quite high among women due to their anatomic and physiologic features such as shorter urethral length, shorter distance from the anus to the urethral meatus, and permissiveness of the vaginal and perineal environments to microbial colonization [1,2]. Urinary tract infections (UTIs) in women represent one of the most common disease entities in both the ambulatory and hospital setting [3]. Nearly 60% of all women will experience at least one UTI in their lifetime leading to billions of dollars of healthcare spending [3].

UTI can be categorized into two: the uncomplicated and complicated UTI. Uncomplicated UTIs also known as uncomplicated cystitis is an infection limited to the bladder or lower urinary tracts. Uncomplicated cystitis is primarily observed in young, healthy women and very rarely requires hospitalization and parenteral antimicrobial therapy. Complicated UTI, on the other hand, is a broad term that encompasses various infections including complicated cystitis, pyelonephritis, and catheter-associated UTI [4]. Migration of the infection towards the kidneys leads to inflammation of the renal parenchyma causing pyelonephritis. Constant inflammation can lead into tubular damage generating interstitial oedema, nephritis and acute kidney injury, thus leading to systemic injuries and damages. Complicated disease most frequently occurs in older adults, those with underlying renal pathology, catheterization, and immune compromising conditions [4].

Oxidative stress relates to increasing intracellular levels of reactive oxygen species (ROS), which is a heterogeneous group of highly reactive ions and molecules generated during univalent reduction of oxygen to water including superoxide anions, hydrogen peroxide, hydroxyl radicals, singlet oxygen and hypochlorous acid [5]. Normally the production of ROS is low and the low levels of ROS are necessary for several biological processes, including intracellular differentiation and cell progression, arrest of growth, apoptosis, immunity and defence against microorganisms. However, oxidative stress caused by increased formation of ROS and/or decreased antioxidant defence may lead to damage of biological macromolecules. This possible damage by ROS can be prevented by the endogenous antioxidant enzymes such as catalase (CAT) and superoxide dismutase (SOD) [6,7]. In bacterial infections oxidative stress arises, at least in part, from altered metabolic pathways and has also been implicated in organ damage and the development of malignancies [8]. The action of bacterial infection in the urinary tract can, thus, trigger the release of free radicals which progresses, thus leading to imbalance between oxidative radicals and antioxidants, hence oxidative changes occur.

Lipid peroxidation is one of the most important expression of oxidative stress induced by ROS [7]. The oxidized low density lipoprotein (Ox-LDL) is an indicator of lipid peroxidation. The Ox-LDL is elevated in oxidative stress associated with a variety of pathological conditions such as arteriosclerosis, inflammation, chronic renal failure, acute myocardial and cerebral infarctions, etc [9]. Hence, Ox-LDL is likely to be pronounced

in cases of complicated UTI such as pelvic inflammatory disease and pyelonephritis. A build up of oxidative stress and lipid peroxidation may lead to chronic systemic injuries which is secondary to urinary tract infections. This study investigated the levels of Ox-LDL and SOD which could serve as a prognostic tool to address the susceptibility or predisposition of women with urinary tract infection to some systemic disease conditions. There is paucity of literature addressing oxidative changes in UTIs, hence the need for the study.

2. METHODS

2.1 Research Design

This research was carried out on women with urinary tract infection registered with the University of Nigeria Teaching Hospital (UNTH), Ituku-Ozalla, Enugu State, Nigeria. The subjects were chosen using a simple random sampling technique.

2.2 Study Population

About forty (40) women diagnosed of urinary tract infections (UTIs) at the University of Nigeria Teaching Hospital (UNTH) were recruited for the study. These patients were further divided into two groups: Group One comprises of twenty (20) patients with Gram-negative urinary tract infections and Group Two comprises of twenty (20) patients with Gram-positive urinary tract infections. Also twenty (20) women who were not suffering from UTIs and who were not registered with UNTH were used as a control group for this study.

2.3 Duration of Study

This research was carried out from July to September, 2021.

2.4 Inclusion and Exclusion Criteria

The inclusion criteria used for this study includes women diagnosed with Gram positive or Gram negative bacterial related UTIs, women between the ages of 18 and 55 years and women who are not on antimicrobial therapy already. However, women below 18 years and above 55 years of age, women with viral or fungal related UTIs, women on antimicrobial therapy and women with hypertension, cardiomyopathy, diabetes mellitus, tuberculosis, obesity and HIV/AIDS were excluded from the study. Moreover, women with no sort of UTIs, be it of bacterial, viral or fungal

origin, between 18 – 55 years, not on any antimicrobial therapy and are not admitted to the hospital, were recruited as the control group for this study.

2.5 Specimen Collection

The serum level of oxidized low density lipoprotein (Ox-LDL) and serum level of superoxide dismutase (SOD) were estimated in each of the blood specimen collected. About 5ml of venous blood was collected aseptically by a venepuncture from each consenting subject and dispensed into a plain venoject bottle and transported to the laboratory within 30 minutes. The blood specimens were centrifuged at 500 x g for 5 minutes and the serum were transferred into labelled sterile cryovials and stored at -20°C until ready for analysis. Each specimen was labelled with the subject's initials and laboratory identification number.

2.6 Method of Sample Processing

The Enzyme-Linked Immunosorbent Assay (ELISA) technique was used to determine the concentration of human ox-LDL and SOD in the samples. The ELISA kits were procured from Elabscience kit. The tests were carried out according to manufacturer's instructions. The optical densities (OD) were read at 450nm wavelength using a microtitre plate reader and reported as concentrations of ox-LDL and SOD in pg/ml.

2.7 Data Analysis

The data were analysed using analysis of variance (ANOVA). Statistical analysis of difference was estimated using ANOVA and correlation and values were reported as Mean±SEM. The statistical package for social sciences (SPSS) version 18 will be used. A value of $p < 0.05$ is considered significant.

3. RESULTS

Patients diagnosed with UTI of either Gram-positive or Gram-negative bacteria origin showed decreased Ox-LDL mean difference when compared to control, but did not show statistically significant difference ($P > 0.05$). There was, however, significant ($P < 0.05$) mean difference in SOD among control group compared to both Gram positive UTI patients and Gram negative UTI patients respectively.

Table 1. Percentage of causative organisms in the women diagnosed with UTI

	Isolated Organism	Number of Isolates	Percentage %
Gram +ve	<i>Staphylococcus aureus</i>	14	70
	<i>Streptococcus</i> spp.	4	20
	<i>Enterococcus faecalis</i>	2	10
Gram -ve	<i>Klebsiella</i> spp.	2	10
	<i>Escherichia coli</i>	16	80
	<i>Proteus</i> spp.	2	10

Table 2. Oxidized Low Density Lipoprotein and Superoxide Dismutase values among Gram positive UTI patients, Gram negative UTI patients and control group

	Control n=20	Gram +ve UTI n=20	Gram -ve UTI n=20	P-value
Ox-LDL (pg/ml)	3900.05±292.89	3854.05±49.52	3798.44±96.35	0.06
SOD (pg/ml)	4064.99±344.66	3790.44±600.62	3651.01±120.21	0.00

4. DISCUSSION

Reactive oxygen species such as superoxide radicals are thought to underlie the pathogenesis of various diseases. Oxidative stress is seen as a situation where there is increased formation of ROS and/or decreased antioxidant defence, an important feature of many diseases and in severe cases, uncontrolled lipid peroxidation is induced [6,10]. To check for possible oxidative stress in UTIs in this study, the SOD, an endogenous antioxidant enzyme, and Ox-LDL, a marker for lipid peroxidation were quantitatively measured. The study showed a significantly decreased mean value of SOD among the test groups compared to the control group. However, lower SOD mean value was seen in Gram negative bacteria UTI infections which are the commonest causative agents of UTI, than in UTI caused by Gram positive bacteria. In the studies of Kurutas et al. [7] on the effects of oxidative stress in urinary tract infection, two antioxidant enzymes measured, catalase (CAT) and SOD, were significantly decreased in the urine samples of patients with UTI compared to negative urine culture. Also a study by Ciragil et al. [10] on the effects of oxidative stress in urinary tract infection during pregnancy shows a reduced CAT and SOD values in pregnant women with UTI compared to pregnant women without UTI and non pregnant women.

Ox-LDL is an indicator of lipid peroxidation, a product of severe oxidative stress. In this study, there was a decrease in the mean value of Ox-LDL in patients with UTI compared to the control. In a case of oxidative stress, it is expected to have higher Ox-LDL in patients with UTI than the control subjects. But given the fact that Ox-LDL is seen in conditions such as atherosclerosis,

inflammation, chronic renal failure, acute myocardial and cerebral infarctions, etc [9], its mean value should increase mainly in cases of complicated UTI with kidney involvements such as pyelonephritis, renal failure, etc. A mean value lower than that of the control values could mean that none of the patient is suffering from a complicated form of UTI. Therefore one can deduce that none of the patients recruited for this study had a complicated form of UTI. However, the studies of Ciragil et al. [10] and Kurutas et al. [7] revealed significant lipid peroxidation as shown by increased levels of malondialdehyde (MDA), another marker for lipid peroxidation, among UTI patients.

5. CONCLUSION

Oxidative stress can underlie urinary tract infection as seen in this study, and also lipid peroxidation in many serious cases, which can further diminish the levels of antioxidant enzymes. A build-up of ROS and lipid peroxidation could lead to systemic injuries secondary to UTIs and so checking for and treating oxidative stress while handling cases of UTI by clinicians in addition to antimicrobial regimen should be seriously taken into cognizance..

CONSENT

It is not applicable.

ETHICAL CONSIDERATION

The following ethical considerations were undertaken during the course of this study: Ethical approval was sought and obtained from the Health Research and Ethics committee

(HREC) of the University of Nigeria Teaching Hospital, Ituku–Ozalla, Enugu State, Nigeria. Each participant signed a voluntary informed consent form prior to participation in the study. Participants' privacy and confidentiality were maintained and only data obtained from this study were presented.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Priya BB. A review of urinary tract infection (UTI). World Journal of Pharmaceutical Research. 2019;9(1):602–25.
2. Mclellan LK, Hunstad DA. Urinary tract infection: Pathogenesis and outlook. Trends in Molecular Medicine. 2016; 22(11):94–57.
3. Matulay JT, Mlynarczyk CM, Cooper KL. Urinary tract infections in women: Pathogenesis, diagnosis and management. Current Bladder Dysfunction Reports. 2016;11:53–60.
4. Steiger SN, Comito RR, Nicolau DP. Clinical and Economic Implications of Urinary Tract Infections. Expert Review of Pharmacoeconomics & Outcomes Research. 2017;17(4):377–83.
5. Miyata Y, Matsuo T, Mitsunari K, Asai A, Ohba K. A review of oxidative stress and urinary dysfunction caused by bladder outlet obstruction and treatments using antioxidants. Antioxidants. 2019;8(5):132.
6. Bratovcic A. Antioxidant enzymes and their role in preventing cell damage. Acta Scientific Nutritional Health. 2020;4(3):1–7.
7. Kurutas EB, Ciragil P, Gul M, Kilinc M. The effects of oxidative stress in urinary tract infection. Mediators of Inflammation. 2005;4:242–4.
8. Ivanov AV, Bartosch B, Isaguliant MG. Oxidative stress in infection and consequent disease. Oxidative Medicine and Cellular longevity; 2017, Article ID: 3496043.
9. Falih IQ, Alobeady MAH, Banoon SR, Saleh MY. Role of oxidized low-density lipoprotein in human diseases: A review. Journal of Chemical Health Risks. 2021;11:71–83.
10. Ciragil P, Kurutas EB, Gul M, Kilinc M, Aral M, Guven A. The effects of oxidative stress in urinary tract infection during pregnancy. Mediators of Inflammation. 2005;5:309–11.

© 2024 Uzoukwu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/102977>