



Study of Physiological Growth Indices and Yield of Chickpea (*Cicer arietinum* L.) to Soil and Foliar Application Through Integrated Nutrient Management Practices

Sri Abisankar M ^{a*}, Augustine R ^a, Isaac Manuel R ^a,
Balaganesh B ^b and Dinesh Kumar ^c

^a Division of Agronomy, School of Agricultural Science, Karunya Institute of Technology and Sciences, Coimbatore, India.

^b Division of Soil Science and Agricultural Chemistry, School of Agricultural Science, Karunya Institute of Technology and Sciences, Coimbatore, India.

^c Division of Agricultural Statistics, School of Agricultural Science, Karunya Institute of Technology and Sciences, Coimbatore, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJSSPN/2024/v10i2275

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116061>

Original Research Article

Received: 10/02/2024
Accepted: 15/04/2024
Published: 16/04/2024

ABSTRACT

An investigation was conducted in Rabi season of 2023-24 at the instructional farm of Karunya Institute of Technology and Sciences, Coimbatore, to evaluate the performance of physiological indices on chickpeas under soil and foliar application. The experiment was established on a

*Corresponding author: Email: abikabi.avm@gmail.com;

Factorial Randomized Block Design (FRBD). The treatments were soil application of NPK 100 percent (A₁), NPK 100 percent + FYM 10 t ha⁻¹ (A₂), NPK 100 percent + Vermicompost 5 t ha⁻¹ (A₃), NPK 75 percent + FYM 10 t ha⁻¹ (A₄) and NPK 75 percent + Vermicompost 5 t ha⁻¹ (A₅) combined with foliar applications, viz., Nano-DAP (B₁) and Nano-urea (B₂). The results of the study revealed that morpho-physiological attributes of chickpeas were statistically increased by the application of soil (NPK 75 percent + Vermicompost 5 t ha⁻¹) combined with foliar (Nano-urea) nutrient management practices resulting in maximum grain yield of 15.64 q ha⁻¹ and stover yield of 30.49 q ha⁻¹. The results of this study underscore the importance of integrated nutrient management practices involving soil and foliar applications in enhancing the physiological indices of chickpeas.

Keywords: Chickpea; nano-DAP; nano-urea; crop growth rate; relative growth rate; chlorophyll index.

1. INTRODUCTION

Chickpea (*Cicer arietinum* L.) is India's second major pulse crop after the common bean (*Phaseolus vulgaris*). Chickpea is grown predominantly under rainfed conditions. It benefits farming systems with a lower carbon footprint due to biological nitrogen fixation to increase soil health [1]. Chickpea is grown in an area of 15.1 million ha (hectares) worldwide, with a total production of 15.8 million tonnes [2]. India and Australia are the top producers, contributing together 73 percent of the total area and production [3]. Pulses are cultivated in an area of 27.9 million hectares in India, with a total production of 23.1 million tonnes. Meanwhile, chickpea is grown in an area of 13.7 million ha with a total production of 10.2 million tonnes with productivity of 1447 kg ha⁻¹ in India and 40.6 lakh hectares of area with 37.67 lakh production and 9.23 lakh productivity in Tamil Nadu [4].

Chickpeas can fix N up to 140 kg ha⁻¹ from the atmosphere and receive 80 percent of their nitrogen (N) needs through symbiotic nitrogen fixation. It leaves a sizable quantity of residual nitrogen for succeeding crops and contributes as organic matter that helps maintain and improve soil health. Plant nutrients are the primary sources for increasing the quality and quantity of chickpeas. Nutrient availability is one of the significant constraints for food production and soil fertility, with unequal utilization of plant nutrients affecting crop growth, development, and yield [5]. The sole application of chemical fertilizers results in good crop production, but it affects soil health and status. Therefore, effective nutrient management can be done by integrating chemical fertilizers with organic manures such as farmyard manure and vermicompost.

Integrated Nutrient Management (INM) plays a crucial role in agricultural sustainability, encompassing the integration of synthetic

fertilizers, organic manure, compost, biofertilizers, and micronutrients. By combining these elements, INM aims to optimize crop yields while minimizing nutrient losses and ensuring agricultural profitability. Among the crop nutrients, nitrogen stands out for its pivotal role in plant growth and development. Consequently, adopting foliar application of nutrients emerges as a more efficient strategy than traditional fertilization methods. Foliar fertilization involves directly spraying or applying liquid or water-soluble fertilizers onto plant leaves, facilitating rapid nutrient absorption by bypassing the soil uptake pathway. This supplementary feeding approach proves particularly beneficial during periods of nutrient deficiencies, stress, or accelerated growth, ensuring timely nutrient delivery to support plant health and productivity. Foliar nutrition minimizes nutrient loss, boosts bioavailability, and economizes crop output by lowering cultivation costs and reducing the quantity of fertilizer applied to crops [6,7].

Nano-urea, a new agricultural input based on nanotechnology, has a particle size of 20 to 50 nm, offering a substantially higher surface area than regular urea prills. Liquid Nano-urea, when directly sprayed onto leaves, enables absorption through stomata, providing crops with a targeted nutrient. Nano-DAP, a white liquid fertilizer that provides phosphorous and nitrogen to plants in a 2.5 :1 ratio, improves crop growth and yields. Nano-DAP has great absorption capacity and easily penetrates plant tissues through stomata when used as a foliar spray [8]. Additional nutrient delivery is a critical aspect in boosting grain production in legumes. The degree of bloom drops impacts chickpea production, contributing features, and yield. The plant's retention of blossoms results in a more significant yield than projected [9]. The present study is aimed to determine the physiological attributes of chickpeas through the combined application of soil and foliar applications of nano-fertilizers.

2. MATERIALS AND METHODS

A field experiment was carried out at Karunya Institute of Technology and Science, Coimbatore, Tamil Nadu to study the response of chickpea (*Cicer arietinum* L.) to soil and foliar application through integrated nutrient management practices under Coimbatore region during *Rabi* season of 2023-24. The climatic condition under Coimbatore district of Tamil Nadu is subtropical. The total rainfall received during the crop-growing period was 128 mm. The weekly maximum and minimum temperature during the experimental period ranged from 21.0°C to 35.1°C and 5.2°C to 15.1°C, respectively. The soil of the experimental plot was sandy clay loam with low organic carbon (0.42 percent), low in available nitrogen (164 kg ha⁻¹), high in available phosphorus (28.5 kg ha⁻¹) and low in available potassium (235 kg ha⁻¹). The soil reaction of the experimental field was alkaline (pH 8.10) with an electrical conductivity of 0.28 dSm⁻¹.

List 1. Treatment details

A Factor (Soil)	Application
(A ₁)	NPK 100 percent
(A ₂)	NPK 100 percent + FYM 10 t ha ⁻¹
(A ₃)	NPK 100 percent + VC 5 t ha ⁻¹
(A ₄)	NPK 75 percent + FYM 10 t ha ⁻¹
(A ₅)	NPK 75 percent + VC 5 t ha ⁻¹
B Factor (Foliar)	Application
(B ₁)	Nano-DAP
(B ₂)	Nano-Urea

*NPK ratio of 25:50:20

The experiment was laid out in factorial randomized block design with soil application of NPK 100 percent (A₁), NPK 100 percent + FYM 10 t ha⁻¹ (A₂), NPK 100 percent + Vermicompost 5 t ha⁻¹ (A₃), NPK 75 percent + FYM 10 t ha⁻¹ (A₄) and NPK 75 percent + Vermicompost 5 t ha⁻¹ (A₅) combined with foliar applications, viz., Nano-DAP (B₁) and Nano-Urea (B₂). The chickpea seeds are sown in well-prepared land by dibbling method with seed rates of 90 kg ha⁻¹. A spacing of 30 × 10 cm was adopted. The crop was produced in an irrigated environment with one pre-planting irrigation applied 7 days before sowing. Hand weeding was done between crops and rows 30 to 45 days after sowing. The crops were harvested manually using a sickle, wrapped into bundles with tags from each plot, and sundried. Threshing procedures were also carried out manually and treatment-wise. The data

collected on various characters studied during the experiment were subjected to statistical analysis in a factorial randomized block design (FRBD). The significance of the difference was tested by the “f” test at a 5 percent level.

2.1 Crop Growth Rate

The crop growth rate (g⁻¹m⁻²day) for each specified stage was calculated using the standard formula given by Radford [10] below:

$$CGR = \frac{W_2 - W_1}{p(T_2 - T_1)}$$

Where,

W₂ = Dry weight of crop plant at the time interval T₂

W₁ = Dry weight of crop plant at the time interval T₁

p = ground area occupied by the plant in m²

2.2 Relative Growth Rate

The relative growth rate (g⁻¹g⁻¹day) for each observational stage was worked out by substituting the corresponding dry matter accumulation values of that very stage in the formula was given by Radford [10] under:

$$RGR = \frac{\text{Log}_e W_2 - \text{Log}_e W_1}{T_2 - T_1}$$

Where,

W₂ and W₁ are the dry matter of plants at the time of T₂ and T₁

2.3 Chlorophyll Rate

The chlorophyll content of the crop was measured using an At LEAF Digital Chlorophyll Meter, which was very easy to handle, and the values were recorded quickly and perfect SPAD (Soil Plant Analysis Development) value were maintained properly.

3. RESULTS AND DISCUSSION

3.1 Crop Growth Rate

The result showed that the effect of combined application of soil and foliar application with NPK 75 percent + Vermicompost along with Nano-Urea (A₅B₂) resulted in a higher crop growth rate

60 DAS to at harvest $2.36 \text{ g}^{-1}\text{m}^{-2}\text{day}^{-1}$. The lower crop growth rate was observed in the combination of 100 percent NPK with Nano-DAP $0.81 \text{ g}^{-1}\text{m}^{-2} \text{ day}^{-1}$ (A₁B₁) (Table 1). The growing trend of CGR might be attributed to increased photosynthetic activity, and a favourable response of CGR leads to an increased plant population. Similar results were also reported by Edwards *et al.*, [11] in maize (Fig. 1). CGR decreases after harvest owing to leaf senescence and a drop in leaf area index.

3.2 Relative Growth Rate

The result showed that the effect of combined treatment of soil and foliar application with NPK 75 percent + Vermicompost along with Nano-Urea (A₅B₂) resulted in a higher relative growth rate at 65 DAS- At harvest ($0.0230 \text{ g}^{-1}\text{g}^{-1}\text{day}$), whereas the lower relative growth rate was observed in the combination of (A₁B₁) 100 percent NPK with

Nano-DAP $0.0200 \text{ g}^{-1}\text{g}^{-1}\text{day}$ (Table 1). The combined application of NPK 75 percent + Vermicompost with Nano-Urea greatly influences relative growth parameters, constituting a multifaceted approach to augment plant growth and productivity. The treatment enhances nutrient availability, soil structure, and stress tolerance, thereby eliciting profound physiological responses, notably reflected in the RGR of plants. With more macronutrients, micronutrients, and organic matter, the treatment optimizes metabolic processes, energy utilization, and biomass accumulation. Uniformly released, the Nano-Urea sustains nutrient availability throughout the growth cycle. A higher supply of nitrogen boosts plant growth and enhances physiological activities, thereby amplifying the production of growth and yield components (Fig. 2). These result findings were in close agreement with the findings of Saitheja *et al.*, [12] on the green gram, Omran *et al.*, [13] on mung bean.




Fig. 1. Effect of Nano-fertilizers on crop growth rate




Fig. 2. Effect of nano-fertilizers on relative growth rate

Table 1. Effect of nano-fertilizer on crop physiological growth indices

Treatments	Crop growth rate (g ⁻¹ m ⁻² day ⁻¹)			Relative growth rate (g ⁻¹ g ⁻¹ day ⁻¹)			Chlorophyll rate			Crop yield (q ha ⁻¹)	
	30-45 DAS	45- 60 DAS	60 DAS- Harvest	30-45 DAS	45- 60 DAS	60 DAS- Harvest	At vegetative	At initial flowering	At peak flowering	Grain yield	Stover yield
Soil Application											
NPK 100 percent	0.38	0.84	1.21	0.0385	0.0425	0.0213	13.15	31.75	35.40	8.08	15.80
NPK 100 percent + FYM 10 t ha ⁻¹	0.43	0.94	1.35	0.0395	0.0430	0.0215	14.99	32.30	36.40	8.20	16.13
NPK 100 percent + VC 5 t ha ⁻¹	0.49	1.09	1.56	0.0395	0.0435	0.0218	15.90	33.40	36.96	9.93	19.63
NPK 75 percent + FYM 10 t ha ⁻¹	0.56	1.22	1.77	0.0400	0.0435	0.0218	16.00	35.45	37.85	11.59	22.70
NPK 75 percent + VC 5 t ha ⁻¹	0.59	1.29	1.87	0.0400	0.0435	0.0218	16.60	36.77	40.90	11.89	23.24
S.E (d) ±	0.01	0.03	0.05	0.0002	0.0001	0.0001	0.42	0.23	0.38	0.30	0.57
CD (p=0.05)	0.03	0.07	0.10	0.0004	0.0002	0.0001	0.89	0.49	0.81	0.64	1.21
Foliar Application											
Nano-DAP	0.32	0.63	1.03	0.0364	0.0408	0.0204	10.28	28.29	481.29	7.39	14.55
Nano-Urea	0.66	1.52	2.08	0.0426	0.0456	0.0228	20.38	39.58	643.74	12.46	24.45
S.E.(d) ±	0.02	0.05	0.07	0.0003	0.0002	0.0001	0.67	0.37	0.61	0.48	0.91
CD (p=0.05)	0.05	0.12	0.15	0.0006	0.0004	0.0002	1.41	0.77	1.28	1.02	1.91
Interaction (AxB)											
S.E. ±	0.03	0.08	0.10	0.0004	0.0003	0.0001	0.95	0.52	0.86	0.68	1.29
CD (p=0.05)	0.07	0.16	0.22	0.0009	0.0005	0.0003	1.99	1.08	1.81	1.44	2.71




Fig. 3. Effect of nano-fertilizers on chlorophyll rate

3.3 Chlorophyll Rate

The result showed that the effect of combined treatment of soil and foliar application with NPK 75 percent + Vermicompost along with Nano-Urea (A_5B_2) resulted in higher during peak flowering stage at 65 DAS - At harvest (0.0230), whereas the lower chlorophyll content at peak flowering stage was observed in the combination of 100 percent NPK with Nano-DAP (A_1B_1 - 0.0200) (Table 1). The foliar application of water-soluble fertilizers accelerated crop growth during the early stages and gradually slowed as the crop matured. This phenomenon may be attributed to the crop's rapid absorption of macro and micronutrients. Increased total chlorophyll and enzyme activities were observed, leading to enhanced photosynthesis. Due to the heightened leaf thickness, there is a likelihood of increased chlorophyll density within the leaves, thereby potentially sustaining more efficient photosynthesis (Fig. 2). These are similar findings collaborated with Kashiwagi *et al.*, [14] and Kumar *et al.*, [15] on sesame.

3.4 Crop Yield

The combined application of treatment NPK 75 percent + Vermicompost along with Nano-Urea (A_5B_2) recorded the maximum grain yield of (15.64 q ha^{-1}) and stover yield of (30.49 q ha^{-1}) resulting in higher quantity of seeds and pods per plant, that increased the crop yield. This increase might be due to the spraying of nano nitrogen during flowering phases, which provides nutrients that promote the performance of the photosynthetic process Anna Joy and Vandha,

[16]. It favourably influences increasing the yield, or the cause is that plants have more pods, which boosted total seed yield Hossain *et al.*, [17]. When applied externally, Nano urea improves photosynthetic capability and reduces the rate at which chlorophyll and leaf nitrogen are lost, hence boosting seed production.

4. CONCLUSION

The findings of this study indicate that combined application of soil and foliar NPK 75 percent, Vermicompost, and Nano-Urea exhibited remarkable efficacy in enhancing chickpea growth and productivity. The combination effect of soil and foliar applications was significant. This increase in production resulted from increased chickpea crop growth and development with the application of organic and inorganic fertilizers. It might be due to the higher nutrient availability during crop growth, which eventually boosted growth. The treatment with soil application of NPK 75 percent + VC 5 t ha^{-1} with foliar application of Nano-Urea led to significantly higher crop growth rates, relative growth rates, and chlorophyll rates compared to other combinations because of its high solubility and rapid absorption characteristics by plant leaves than Nano-DAP, as phosphorus tends to be less mobile in plant tissues. These improvements can be attributed to the multifaceted approach of nutrient supply, enhanced soil structure, and stress tolerance. The sustained release of nutrients from nano played a pivotal role in optimizing metabolic processes and biomass accumulation throughout the growth cycle.

ACKNOWLEDGEMENT

The authors are grateful to the Division of Agronomy, School of Agricultural Sciences (SAS), Karunya Institute of Technology and Sciences (KITS), Coimbatore, Tamil Nadu – 641114.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bohra A, Pandey MK, Jha UC, Singh B, Singh IP, Datta D, Varshney RK. Genomics-assisted breeding in four major pulse crops of developing countries: present status and prospects. *Theoretical and Applied Genetics*. 2014;127:1263-1291.
2. Faostat, Food and Agricultural Organization of the United Nations, [faostat.fao.org](http://www.fao.org/statisticaldatabases/faostat); 2021.
3. Gaur PM, Samineni S, Thudi M, Tripathi S, Sajja SB, Jayalakshmi V, Dixit GP. Integrated breeding approaches for improving drought and heat adaptation in chickpea (*Cicer arietinum* L.). *Plant Breeding*. 2019; 138(4):389-400.
4. Directorate of Economics and Statistics, *Agricultural Statistics at a Glance*; 2023. Available:<http://desagri.gov.in/>
5. Siddiqui SN, Umar S, Husen A, Iqbal M. Effect of phosphorus on plant growth and nutrient accumulation in a high and a low zinc accumulating chickpea genotypes. *Annals of Phytomedicine*. 2015;4(2):102-105.
6. Srivastava A, Singh R. Effect of Nitrogen and Foliar Spray of Urea and Nano Urea on Growth and Yield of Rabi Maize (*Zea mays* L.). *International Journal of Plant & Soil Science*. 2023;35(18):2037-2044.
7. Kumar A, Ram H, Kumar S, Kumar R, Yadav A, Gairola A, Sharma T. A Comprehensive Review of Nano-Urea vs. Conventional Urea. *International Journal of Plant & Soil Science*. 2023; 35(23):32-40.
8. Maloth A, Thatikunta R, Parida BK, Naik DS, Varma N. Evaluation of Nano-DAP on Plant Growth, Enzymatic Activity and Yield in Paddy (*Oryza sativa* L.). *International Journal of Environment and Climate Change*. 2024;14(1): 890-897.
9. Kamaleshwaran R, Karthiga S. Effect of foliar nutrition on yield and growth parameters of greengram in coastal area of Tamil Nadu (*Vigna radiata*) CV. Vamban 2. *International multidisciplinary e-Magazine*. 2021;1(3): 22-27.
10. Radford PJ. Growth analysis formulae-their use and abuse 1. *Crop science*. 1967;7(3): 171-175.
11. Edwards JT, Purcell LC, Vories ED. Light interception and yield potential of short-season maize (*Zea mays* L.) hybrids in the Midsouth. *Agronomy Journal*. 2005; 97(1):225-234.
12. Saitheja V, Senthivelu M, Prabukumar G, Prasad V. Maximizing the Productivity and Profitability of Summer Irrigated Greengram (*Vigna radiata* L.) by Combining Basal Nitrogen Dose and Foliar Nutrition of Nano and Normal Urea. *International Journal of Plant and Soil Science*. 2022;34(22):109-116.
13. Omran AH, Dass A, Jahish F, Dhar S, Choudhary AK, Rajanna GA. Response of mungbean (*Vigna radiata* L.) to phosphorus and nitrogen application in Kandahar region of Afghanistan. *Annals of Agricultural Research*. 2018;39(1).
14. Kashiwagi J, Upadhyaya HD, Krishnamurthy L. Significance and genetic diversity of SPAD chlorophyll meter reading in chickpea germplasm in the semi-arid environments. *Journal of food legumes*. 2010;23(2):99-105.
15. Kumar A, Singh SN, Khan MA. To know the effect of foliar application of thiourea and potassium nitrate on physiological growth at different stages of sesame (*Sesame indicum* L.). *Journal of Pharmacognosy and Phytochemistry*. 2018;7(2):2156-2158.
16. Anna Joy and Vandna Chhabr. Effect of Application of Nano-nutrients on Crop Growth and Seed Yield in Black Gram (*Vigna mungo*) Ecology, Environment and Conservation. 2023;29.

17. Hossain ME, Begum S, Hoque MM, Islam MM, Shahed M. Foliar Application of Nano Urea on Growth and Yield of Binamash-3 (*Vigna mungo* L.). Journal of Agroforestry and Environment. 2023;16 (2):131-133.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/116061>