



## **Cavoatrial Thrombus Revealing a Renal Cell Carcinoma**

**A. El Adaoui <sup>a\*</sup>, I. Tlohi <sup>a</sup>, H. Bendahou <sup>a</sup>, R. Habbal <sup>a</sup>, H. Lachhab <sup>b</sup>  
and M. Graioud <sup>b</sup>**

<sup>a</sup> *Cardiology Department, Ibn Rochd University Hospital, Casablanca, Morocco.*

<sup>b</sup> *Urology Department, Ibn Rochd University Hospital, Casablanca, Morocco.*

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

#### Editor(s):

(1) Dr. Prabakaran Nagarajan, College of Medicine, The Ohio State University, USA.

#### Reviewers:

(1) Mirela Gliga, Romania.

(2) Kajal Patel, India.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/80347>

**Received 15 October 2021**

**Accepted 27 December 2021**

**Published 29 December 2021**

### **Case Study**

### **ABSTRACT**

Renal carcinoma is a tumor with high metastatic risk, compromising its prognosis, often fatal. Involvement in the inferior vena cava is present in 15-20% of metastatic locations with the possibility of intracavitary cardiac extension in 5%.

The circumstance of discovery of this tumor are variable, and often discovered by chance on extra-renal explorations.

The right cavo-atrial localization as a sign of discovery has been rarely described in the literature. We report the observation of a young patient who consulted for chest pain, subsequently demonstrating a metastatic clear cell carcinoma to the cavo-atrial system.

*Keywords: Cavo-atrial thrombus; renal carcinoma; insidious thrombus.*

### **ABBREVIATIONS**

*CT : Computed Tomography*

*ECG : Electrocardiogram*

*IVC : Inferior Vena Cava*

*MRI : Magnetic Resonance Imaging*

*RCC : Renal Cell Carcinoma*

*TTE : Transthoracic Echocardiography*

\*Corresponding author: E-mail: [Eladaoui.abdelmajid@gmail.com](mailto:Eladaoui.abdelmajid@gmail.com), [abdel2web@hotmail.fr](mailto:abdel2web@hotmail.fr);

## 1. INTRODUCTION

Renal carcinoma is the most common neoplastic entity of renal tumors, often occurring in the sixth decade of life, considered to have a poor prognosis due to its predominant metastatic potential.

Venous extension with thromboembolic complications are special features of renal cell carcinoma, compromising his prognosis [1].

This extension occurs in 10% of patients, with involvement of the right atrium in 1% of cases [2].

We present here the case of a renal carcinoma, discovered in the complicated stage of right cavo-atrial thrombus.

## 2. CASE REPORT

-44-year-old youngster, with no particular history, presented to the emergency department for abdominal distension that had gradually set in for 2 months, complicated by dyspnea at rest with edema of the lower limbs.

-Clinical examination found a patient with dyspnea at rest, with abdominal tenderness, the arterial pressure was 130/76 mmHg, a heart rate at 104 bpm, abdominal collateral circulation, hepatomegaly, and swollen lower limbs.


-The ECG was unremarkable, the transthoracic echocardiography found an aspect of serpiginous

thrombus encumbering the inferior vena cava and arriving intracavitary at the level of the right atrium, without involvement of the tricuspid valve (Fig. 1) and extending upstream to the level of the unilateral primitive iliac and common femoral veins (Fig. 2).


The biological assessment found anemia at 5 g / dL, a slight deterioration of renal function with creatinemia at 15 mg / L, and remained stable after rehydration after the CT imaging; the cytobacteriological examination of the urine showed microscopic hematuria.

The CT imaging objectified: a hypodense endoluminal material over the entire length of the IVC extended downstream of the right atrium, with large left renal mass associated with a magma of lumbo-aortic lymphadenopathy, and diffuse peritoneal and retro-peritoneal infiltration, associated with multiple hepatic masses invading the contiguity of the IVC and staged vertebral osteolytic lesions; we concluded a suspicion of a left renal tumor complicated by hepatic, peritoneal and bones metastases, with invasion of the cavo-atrial system (Fig. 3).


The patient received a blood transfusion and put on anticoagulation; after biological stabilization, he subsequently underwent a left nephrectomy with cavotomy and thrombectomy, histological examination confirmed a clear cell renal carcinoma (Figs. 4-5).



**Fig. 1. A) short-axis TTE image: Mobile thrombotic masses within the right atrium (B) subcostal image: serpiginous thrombus along the inferior vena cava**




**Fig. 2.** Transverse vascular ultrasound image of the primary iliac vessels: endoluminal material encumbering the unilaterals primary iliac and common femoral vein



**Fig. 3.** Thoraco-abdominal CT images: A: left renal mass with diffuse peritoneal infiltration ;B multiple fairly well-limited hepatic masses of secondary origin ; C: endoluminal material at the level of the Inferior vena cava; D: right intra-atrium thrombus



**Fig. 4. Left nephrectomy surgery piece**



**Fig. 5. Histopathological study showing a clear cell renal cell carcinoma**

### 3. DISCUSSION

-Our observation highlights a form of fortuitous discovery of renal carcinoma, based on thromboembolic complications.

-The typical triad of kidney tumors: flank pain, mass and hematuria was not present in our

patient. and the diagnosis was oriented through the discovery of the thrombotic images on the echocardiography, the differential diagnosis was initially made with a cardiac tumor (myxoma), a thrombus of the eustachi valve. The serpiginous extension at the level of the IVC and the venous network upstream directed to search for a primary localization.

-In the literature, similar cases have been reported .Most of the cases have described insidious discoveries on request for imaging, sometimes finding metastases from the outset as is the case with us [3-4-5].

-Recourse to multimodal imaging is often essential to assess the extent, complications and guide therapeutic management, renal ultrasound is the first accessible examination, followed by abdominal-thoracic CT, MRI as well as cardiac imaging [6].

-The renal cell carcinoma (RCC), often takes extension at the level of the right cardiac chambers, the precision of the upper margin is important to guide the therapeutic management [6].


RCC is described on 4 levels according to this upper edge:

- \* Level I: the tumor thrombus extends from the renal vein into the IVC, but still infrahepatic level.
- \*Level II: the tumor thrombus extends to the subhepatic IVC
- \*Level III: the tumor thrombus extends into the intrahepatic IVC
- \*Level IV: the tumor extends to the right atrium

- The surgical management in patients with renal cell carcinoma is quite complex, to find a fairly safe result.

Combined surgical intake with the use of cardiopulmonary bypass may be necessary allowing complete resection of the tumor [7-8-9].

-Long-term postoperative survival has been described in studies in spite of the invasive nature of the intracardiac extension [10].



**Fig. 6. Classification of the renal cell carcinoma according to tumor and tumor thrombus level**

#### 4. CONCLUSION

The discovery of extensive cavo-atrial thrombus should lead to a search for a renal or hepatic neoplastic origin.

The use of imaging techniques combining echocardiography, CT and MRI is essential to determine the extent of thrombosis and tumor.

Multidisciplinary urological, vascular and cardiac management is essential to guide safe tumor resection.

#### CONSENT AND ETHICAL APPROVAL

As per university standard guideline, participant consent and ethical approval have been collected and preserved by the authors

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Quencer KB, Friedman T, Sheth R, Oklu R. Tumor thrombus: incidence, imaging, prognosis and treatment. *Cardiovasc Diagn Ther.* 2017;7(Suppl 3):S165-S177.  
DOI: 10.21037 / cdt.2017.09.16  
PMID: 29399520  
PMCID: PMC5778532
2. Schimmer C, Hillig F, Riedmiller H, Elert O. Surgical treatment of renal cell carcinoma with intravascular extension. *Interact Cardiovasc Thorac Surg.* 2004;3(2):395-7.  
DOI: 10.1016 / j.icvts.2004.02.014  
PMID: 17670271
3. Lu HT, Chong JL, Othman N et al. An uncommon and insidious presentation of renal cell carcinoma with tumor extending into the inferior vena cava and right atrium: a case report. *J Med Case Reports.* 2016;10:109.  
DOI.ORG/10.1186/S13256-016-0888-5
4. Posacioglu H, Ayik MF, Zeytunlu M, Amanvermez D, Engin C, Apaydin AZ. Management of renal cell carcinoma with intracardiac extension. *J Card Surg.* 2008;23(6):754-8.  
DOI: 10.1111/j.1540-8191.2008.00664.x.  
PMID: 19017006.
5. Yoon S, Jeon DW, Yang JY. A huge thumb in the heart. *Heart Asia.* 2013;5: 228.  
DOI: 10.1136/heartasia-2013-010443
6. Kakoti S, Jena R, Sureka SK, Srivastava A, Mandhani A, Singh UP. Experience with management of renal cell carcinoma with inferior vena cava/right atrial tumor thrombus. *Indian J Urol.* 2021;37(3):234-240.  
DOI: 10.4103/iju.IJU\_13\_21  
Epub 2021 Jul 1  
PMID: 34465952  
PMCID: PMC8388349
7. Chiappini B, Savini C, Marinelli G, Suarez SM, Di Eusano M, Fiorani V, Pierangeli A. Cavoatrial tumor thrombus: single-stage surgical approach with profound hypothermia and circulatory arrest, including a review of the literature. *J Thorac Cardiovasc Surg.* 2002;124(4):684-8.  
DOI: 10.1067/mtc.2002.124295  
PMID: 12324725.
8. Dominik J, Moravek P, Zacek P, Vojacek J, Brtko M, Podhola M, et al. Long-term survival after radical surgery for renal cell carcinoma with tumour thrombus extension

- into the right atrium. *BJU Inter.* 2012;111:59–64.  
DOI: 10.3400/avd.cr.11.00067
9. Zisman A, Wieder JA, Pantuck AJ, Chao DH, Dorey F, Said JW, Gitlitz BJ, deKernion JB, Figlin RA, Belldegrun AS. Renal cell carcinoma with tumor thrombus extension: Biology, role of nephrectomy and response to immunotherapy. *J Urol.* 2003;169(3):909-16.  
DOI: 10.1097/01.ju.0000045706.35470.1e.  
PMID: 12576811.
10. Maninder S. Kalkat, Asad Abedin, Stephen Rooney, Alan Doherty, Muzaffar Farouqi, Michael Wallace, Timothy R. Graham, Renal tumours with cavo-atrial extension: Surgical management and outcome, *Interactive Cardio Vascular and Thoracic Surgery.* 2008;7(6):981–985.  
Available: <https://doi.org/10.1510/icvts.2008.180026>

---

© 2021 Adaoui et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/80347>