



Degradation of Emamectin Benzoate in the Aquaculture Pond Environment under Tropical Climatic Condition

Ramesh Babu Rajesh ^a, Ramamoorthy Saraswathy ^{a*},
Satheesha Avunje ^a,
Ramalingam Ananda Raja ^a, Periyamuthu Kumararaja ^a
and Prasanna Kumar Patil ^a

^a ICAR- Central Institute of Brackishwater Aquaculture, 75 Santhome High Road, MRC Nagar, Chennai, Tamil Nadu, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajee/2024/v23i10607>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/123208>

Original Research Article

Received: 09/07/2024

Accepted: 11/09/2024

Published: 18/09/2024

ABSTRACT

Emamectin Benzoate (EMB), is an anti-parasitic drug, commonly used in aquaculture for prevention and control of parasitic infestations. The drug enters the environment through uneaten feed and faeces, ultimately accumulating in aquaculture pond sediment. Present study reports the influence of abiotic factors like, sunlight exposure (2,454 to 117,500 lux), pH, salinity and soil texture on the degradation of EMB in aquaculture pond water and sediment as per Organization for Economic Cooperation and Development (OECD). Results showed that the degradation of EMB in both water

*Corresponding author: E-mail: R.Saraswathy@icar.gov.in;

Cite as: Rajesh, Ramesh Babu, Ramamoorthy Saraswathy, Satheesha Avunje, Ramalingam Ananda Raja, Periyamuthu Kumararaja, and Prasanna Kumar Patil. 2024. "Degradation of Emamectin Benzoate in the Aquaculture Pond Environment under Tropical Climatic Condition". *Asian Journal of Environment & Ecology* 23 (10):38-48. <https://doi.org/10.9734/ajee/2024/v23i10607>.

and sediment followed first-order kinetics, resulting in half-life of 5.6 days in water and 12.4 days in sediment when exposed to natural sunlight. Higher degradation of EMB was observed under alkaline conditions (pH 8.5) and at lower salinity (0.5ppt). EMB degradation is accelerated in loamy sand soil when exposed to sunlight. Results of the study suggest that in countries like India, where ample sunlight exposure is prevalent throughout the year, EMB degradation in aquaculture pond environment occurs at a rapid pace reducing the risk of accumulation in water or sediment.

Keywords: Abiotic factors; aquaculture; degradation; emamectin benzoate; kinetics

1. INTRODUCTION

Seafood plays a significant role in the global food system and is undoubtedly one of the world's most valuable commodities in the global context. The nutritional value, health benefits and food security and sustainability of seafood has been well recognized [1]. The exponential growth of the human population and the increasing per capita consumption of seafood have led to a rising demand for fishery products. Presently, over seven billion people depend on fish as a source of over 15% of their animal protein intake. In economically disadvantaged coastal regions, this dependency can soar to as high as 90% [2,3]. Aquaculture is one of the India's fastest-growing food production sectors with an export of 1.73 million MT, contributing significantly to foreign exchange revenue to the tune of US\$ 8.1 billion in the year 2022–23 [4]

Effective control of diseases in fish is an important aspect for increasing the fish culture to meet the ever-rising demand of aquatic products. There are many parasites in the aquatic ecosystem which attach to the fish and feed on the mucus, blood and skin, therefore causing damage to the fish [5]. Ectoparasitic copepods (Copepoda: *Caligidae* and *Lernaeidae*), isopods (Isopoda: *Cymothoidae*) and brachiurans (Brachiura: *Argulidae*) particularly parasites, pose a substantial threat to the health and productivity of the global aquaculture industry. These parasites are known to cause considerable economic losses, with an estimated annual impact exceeding 1.05 billion to 9.58 billion US\$ [6]. EMB is a semi-synthetic derivative of avermectins, a group of macrocyclic lactones derived from the soil bacterium *Streptomyces avermitilis* [7]. It is primarily used in aquaculture to control sea lice and other parasites in farmed fish, particularly salmonids. The drug is typically administered orally as feed top dressing and is known for its broad-spectrum efficacy, long-lasting effect and was identified and developed as an anti-parasiticide for both

marine and freshwater-reared fish species. It has been recommended by Food and Drug Administration (FDA) and European Medical Agency (EMA) for the standard treatment of 50µg of EMB kg⁻¹ of fish body weight (BW) d⁻¹ for seven consecutive days [8,9].

As EMB is widely used as an anti-parasitic agent in aquaculture, it is crucial to understand its fate and degradation in environment. It is estimated that about 75% of antibiotics/ therapeutics/ drugs which are induced in the feed eventually reach the pond environment [10-12]. EMB reduces the benthic population as well as their composition.

The degradation of EMB in aquaculture environments is reportedly influenced by various abiotic factors such as water salinity, temperature, pH, sunlight exposure and the presence of other organisms [13,14]. Hydrolysis, photolysis and biodegradation are considered as a main degradation pathways involved in the breakdown. Hydrolysis, facilitated by water and influenced by pH, leads to the cleavage of ester bonds in EMB, resulting in the formation of primary and secondary degradation products [15]. Photolysis, driven by sunlight can contribute significantly to the degradation of EMB as the compound is susceptible to degradation when exposed to UV light [16].

The complete degradation of EMB from sediment is a challenging process. Studies [17,18] have shown that EMB undergoes photolysis, which is influenced by soil parameters. The breakdown of EMB in soil largely depends on the aerobic and anaerobic conditions, as well as the soil's characteristics. Studies on environmental degradation of EMB in tropical climate like India are limited. Hence, the present study reports the influence of abiotic factors on the degradation of EMB in tropical aquatic environments. The understanding gained by the study will help in the development of strategies for the efficient use of EMB and reduce the effect on the pond environment.

2. MATERIALS AND METHODS

2.1 Chemicals and Sample Preparation

Emamectin benzoate, 99.3% pure (Analytical standard), powder ((4''R)-4''-Deoxy-4''-(methylamino) avermectin B1 benzoate) with molecular formula of $C_{56}H_{81}NO_{15}$ and molecular weight of 1008.2 g/mol was obtained from Sigma- Aldrich (Milwaukee, WI, USA). All organic solvents (HPLC grade) were purchased from Sigma-Aldrich and Hi-Media and ultrapure water was used in the preparation of the reagents and purity was maintained. EMB is highly soluble in organic solvents; methanol was used for the preparation of 1000ppm aqueous EMB solution. The stock solution was further diluted into 100ppm using ultrapure water and used for the experiments.

The water (fresh and sea water) and soils from aquaculture ponds used in the experiment was

characterized and checked for any possible contamination with EMB by using Liquid Chromatography Mass Spectrometry (LC-MS/MS) before the start of the experiment. The characteristics of water (Table 1) and soil (Table 2).

2.2 Photochemical Experiments

To study the photodegradation of EMB in water, experiments were conducted in duplicate in three different saline waters (0.5, 15 and 25ppt), three different pH (5,7 and 8.5) and under sunlight and dark conditions. In 250 ml transparent glass (sunlight) and amber colour bottle (dark) with a dimension of 70 mm OD by 143mm height was used, a final concentration of 1000 ppb of EMB solution was taken and the bottles were kept in sunlight for a photolytic degradation. A similar setup was kept in dark environment with constant room temperature at 25 ± 1 °C.




Fig. 1. Variations in the light intensity and temperature during the experimental period

Table 1. Water quality parameters at different pH and salinity levels. (0.5, 15, 25 indicates salinities); (A, B, C indicates pH 5, 7, 8.5 respectively)

	0.5A	0.5B	0.5C	15A	15B	15C	25A	25B	25C
pH	5.05	7.01	8.59	5.03	7.05	8.50	5.07	7	8.60
Salinity (ppt)	0.5	0.6	0.5	14	15	15	25	24	25
Carbonate (ppm as CaCO ₃)	0	0	61.2	0	0	0	0	0	122.4
Bicarbonate (ppm as CaCO ₃)	62.22	273.8	286.2	37.3	124.4	24.888	74.7	223.9	323.5
Total alkalinity (ppm as CaCO ₃)	51	224.4	285.6	30.6	102	81.6	61.2	183.6	265.2
Calcium (ppm)	29.73	29.73	33.98	127.45	212.42	212.42	254.90	339.87	254.90
Magnesium (ppm)	15.61	23.42	15.61	624.53	598.51	572.49	1014.87	910.78	936.80
Total hardness (ppm as CaCO ₃)	137.6	169.6	148.4	2862	2968	2862	4770	4558	4452

Table 2. Soil characteristics of different soil textures

Soil texture	Loamy sand	Clay
pH	8.9	8.46
Electrical Conductivity (mS/cm)	3.24	4.15
Organic carbon (%)	0.02	0.24
Available nitrogen (ppm)	38.65	49.1
Available phosphorus (ppm)	23.75	26.5
Sodium (ppm)	2030	430
Potassium (ppm)	179	136
Pore space volume (ml)	5	7
Pore space %	38.46	41.17
Bulk density (g/cc)	1.53	1.17
Particle density	2.5	2
Calcium (ppm)	27.2	40.7
Magnesium (ppm)	46.3	41.1

In another experiment, two different textured soils (loamy sand and clay) were used and water holding capacity was determined to know the amount of EMB aqueous solution needed to get a final wet weight concentration of 1000ppb. Accordingly, EMB solution was added in 25 g of soil and EMB were mixed using a vortex to get a uniform spread. The soil was kept in sunlight and in the dark to study the degradation of EMB and the experiments were conducted in duplication. The water holding capacity of loamy sand and clay was determined as per the standard method [19] and the same was maintained throughout the experiment.

During the experimental period, data on photoperiod, light intensity (lux) and atmospheric temperature (°C) were recorded (Fig. 1). Water and soil samples were collected once in two days and analyzed in LC-MS/MS for EMB concentration.

2.3 Analytical Determination of LC-MS/MS

The quantification of EMB in the water samples was conducted using Liquid Chromatography Mass Spectrometry (LC-MS/MS). A 10 ml water sample was transferred into a centrifuge tube, followed by the addition of 10 ml of acetonitrile. After vortexing the mixture, 10 g of sodium sulphate was introduced and the solution was subjected to centrifugation (8000 RPM). From the solution, 1 ml was taken and diluted to 10 ml with acetonitrile and injected in LC-MS/MS (Agilent, LC 6470, USA). For soil, 5g of sample was weighed and 10ml water and 10ml acetonitrile in the ratio of 1:1 was added and mixed well. To this mixture, 10g of sodium

sulphate was added, vortexed and centrifuged, 1ml of supernatant was taken and made up to 10ml using acetonitrile and injected. For water samples, the Limit of Detection (LOD) was set at 2 µg/L and the Limit of Quantification (LOQ) was determined to be 5 µg/L. In the case of sediment samples, the LOD and LOQ were respectively defined as 5 µg/kg and 10 µg/kg. These thresholds ensure the reliability and precision of EMB quantification in both water and sediment matrices.

The degradation data for EMB were evaluated using the kinetic equations of both first (a) and second order (b).

For the first-order reaction:

$$\text{Reaction rate} = d[C]/dt = -k [C]$$

$$\text{Integration form of first - order kinetics: } C = C_0 * e^{-kt} \quad (a) \quad [20]$$

The reaction rate is in molar/time, and 'k' is the reaction rate coefficient (time⁻¹). When you plot ln [C] against time for a first-order reaction, you get a straight line. The slope of this line corresponds to the rate constant (k) for the reaction.

The half-life (t_{1/2}) was calculated using the formula: t_{1/2}=0.693/k, 'k' was obtained from the slope of the ln (C) vs time graph.

Second-order reaction equation:

$$\text{Reaction rate} = d[C]/dt = -k [C]^2$$

$$\text{Integration form of second-order kinetics: } 1/C = 1/C_0 + kt \quad (b) \quad [20]$$

In these equations, 'C₀' represents the initial EMB concentration, 'C' is the residual EMB concentration at time 't', and 'k' is the rate constant. The further analysis was done using Computer Assisted Kinetic Evaluation (CAKE) software.

3 RESULTS

3.1 Photodegradation of *Emamectin benzoate* in water

The photodegradation of EMB was examined under natural sunlight with an average sunlight intensity of 35,894 lux, a temperature of 33°C and a photoperiod of 12 hrs 30 mins. The average light intensity showed notable variations based on weather conditions. The highest intensity occurred at approximately 2:00 PM during sunny weather. Meanwhile, a constant dark condition was maintained in a separate room with temperature set at 25±1 °C. The data confirmed exponential decay, indicating that the reaction was first-order. The aqueous solution of EMB was highly stable at pH 5 and it was essentially stable in the dark.

In all the saline water (0.5, 15 and 25 ppt), it was observed that at pH 8.5 tends to degrade rapidly in both sunlight and dark conditions (Fig. 4). The drug in freshwater when exposed to sunlight showed a half-life of 3.03, 4.33 and 5.23 days at pH levels of 8.5, 7 and 5 respectively (Fig. 2).

In contrast, degradation under 25ppt followed a little slower degradation (0.0932, 0.0940 & 0.1027^d) under different pH (pH 5, 7 & 8.5) with a half-life period of 6.75, 7.37 and 7.43 days at pH 8.5, 7 and 5. In comparison with dark conditions, it was higher in all pH with a half-life of 34.3, 54.8 and 55.8 days at pH 8.5, 7 and 5 in low salinity (Fig. 5). The results (Table 3) indicates that EMB undergoes photodegradation, especially at high pH (alkaline) and low saline (0.5ppt) showed to have faster degradation. The degradation rate decreased at lower pH (5) and increased significantly at higher pH (8.5), regardless of the salinity. At higher salinity the degradation was less compared with low saline water. Similarly, in pH 8.5 the half-life period was 3.03 and 6.75 days for freshwater and high saline (25ppt) waters respectively, while it was 5.23 and 7.43 days at pH 5 (Fig. 3).




Fig. 2. The effect of pH on the photodegradation of *Emamectin benzoate* under varying salinities




Fig. 3. The effects of salinities on the photodegradation of *Emamectin benzoate* under varying pH

Table 3. Rate constant (k), half-life (D50) and 90% degradation (D90) of *Emamectin benzoate* under natural sunlight in aqueous solutions

	0.5ppt			15ppt			25ppt		
	pH5	pH7	pH8.5	pH5	pH7	pH8.5	pH5	pH7	pH8.5
K ^{-d}	0.1325	0.1602	0.2286	0.1174	0.1229	0.1331	0.0932	0.0940	0.1027
D50 (days)	5.23	4.33	3.03	5.9	5.64	5.21	7.43	7.37	6.75
D90 (days)	17.4	14.4	10.1	19.6	18.7	17.3	24.7	24.5	22.4

Table 4. Rate constant (k), half-life (D50) and 90% degradation (D90) of *emamectin benzoate* under natural sunlight and dark in soils

	Sunlight		Dark	
	Loamy sand	Clay	Loamy sand	Clay
K ^{-d}	0.0559	0.03037	0.02752	0.01564
D50 (days)	12.4	22.8	25.2	44.3
D90 (days)	41.2	75.8	83.7	147

3.2 Photodegradation of *Emamectin benzoate* in soil

Based on EMB concentration in the soil at periodical intervals under different pH and texture as well as under sunlight and dark, the

degradation rate and half-life period was calculated (Table 4). In natural sunlight, the half-life was 12.4 and 22.8 days in loamy sand and clay soil with a rate constant of 0.0559 and 0.0303 k^{-d}. Whereas in the dark condition, the half-life was about 25.2 and 44.3 days with a rate

constant of 0.0275 and 0.0156^{-d} in loamy sand and clay soil respectively. The data fit into first-order kinetics.

4. DISCUSSION

Emamectin benzoate has been successfully used in farmed fish worldwide as an effective antiparasitic compound. Data on the application and degradation of the compound in tropical climates is scanty and therefore, it is essential to generate data for better environmental practices and safety. In this study, the fate and degradation of EMB in water and sediment under Indian tropical conditions were evaluated. Hydrolysis, photolysis and biodegradation are the main

degradation pathways involved in the breakdown of EMB [21]. The photolysis of the aqueous EMB solution resulted in a notable degradation of its active ingredients, in contrast to the stability observed under dark conditions. The chemical bonds found within the drug, including ether and ester linkages, exhibit susceptibility to photodegradation when exposed to UV light. This photodegradation process can generate numerous by-products, as documented by earlier studies [22-25]. Hydrolysis, facilitated by water and influenced by pH, leads to the cleavage of ester bonds in EMB, resulting in the formation of primary and secondary degradation products [15].




Fig. 4. Comparative degradation of EMB in sunlight vs dark under various salinities




Fig. 5. Comparative degradation of EMB in sunlight vs dark under various pH

4.1 EMB Degradation in Water

In this study, the photolytic degradation rate was more than 10 times as compared with the dark. Earlier studies reported faster photolysis of EMB under natural sunlight compared to artificial light sources [16] as sunlight contains ultraviolet (UV) radiation, particularly in the UVA and UVB ranges, which can initiate photolysis reactions. Photolysis involves the breaking of chemical bonds in EMB by absorbing UV light energy [26]. Some substances present in the environment, such as certain organic matter or metals, can act as sensitizers, which can absorb light energy and transfer to EMB thereby enhancing the degradation process [16]. Photolysis and sensitization reactions can additionally stimulate the production of highly reactive oxygen species, including singlet oxygen, hydroxyl radicals, and superoxide radicals. These reactive species play a pivotal role in driving the degradation of the compound [27].

EMB has varied water solubility under different salinities and is very poor in seawater with only 5.5 mg/l solubility. In this experiment, the degradation was slower with increasing salinity of water with a half-life of 4.19, 5.61 and 7.18 days under 0.5, 15 and 25 ppt salinity respectively. This may be due to the higher concentration of minerals like calcium, magnesium, sodium and potassium in high-saline water. The presence of salts in seawater acts as stabilizers, interacting with the chemical structure of the compound, making it more resistant to degradation by various abiotic factors.

The solubility of EMB changes significantly with the pH of water with a solubility rate of 320 ± 30 mg/L under pH 5, 24 ± 2 mg/L under pH 7 and only 0.1 ± 0.1 mg/L under pH 9. The current study showed that the EMB decay rate is faster in higher pH, it follows as $\text{pH } 8.5 > 7 > 5$ in both photolytic and dark conditions. In a similar study [28], photodegradation of EMB in solution exhibited varying half-life periods. Specifically, EMB had a half-life of 22 days in a pH-buffered solution with a pH of 7 and 7 days in natural pond water (0 ppt). It was reported that EMB was stable at pH 5.2, 6.2, 7.2 and 8.0 at 25°C, whereas at pH 9 the compound breaks down with a half-life of 19.5 weeks under sterile buffered aqueous solution [29]. Therefore, the study shows that as the salinity of the water increases the degradation rate increases with respect to its pH or photolytic conditions. EMB undergoes photolysis and experiences

degradation within the water column at depths where light can reach [22].

4.2 EMB Degradation in Soil

Sunlight was more effective in the degradation of EMB irrespective of the soil texture. The rate of degradation was faster (doubled) under sunlight than in dark conditions. The drug reportedly shows almost no degradation in soil under dark conditions [17]. Between the soil textures, the drug tended to degrade slower in clay soil than in loamy sand under both sunlight and dark. It may be due to large surface area, high cation exchange capacity and overall negative charge, which draws and holds positively charged molecules through electrostatic interaction. Higher organic matter in the clay soil contains various substances, including humic and fulvic acids, which can be complex with EMB forming a stable complex. In addition to this, temperature, moisture and aerobic and anaerobic conditions influences the degradation rate in soil [18].

5. CONCLUSION

In brackishwater and freshwater fish farming, there is a high risk of ectoparasite and endoparasite infections. *Emamectin benzoate* is frequently used to treat and prevent these infections. It's crucial to assess how EMB degrades in various abiotic conditions within the pond environment. The degradation rate of anti-parasitic agent, EMB in both water and soil followed the first-order kinetics and the degradation was faster in low saline water with alkaline pH when exposed to sunlight. Similarly, the degradation was faster under sunlight in the light textured soil than heavy texture soil. The study revealed that under tropical conditions, the risk of EMB accumulation in water and soil is minimal.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscript.

ACKNOWLEDGEMENTS

The authors are grateful to Indian Council of Agricultural Research (ICAR), Ministry of Agriculture and Farmers' Welfare, Government of India for providing financial support through All

India Network Project on Fish Health (AINP_FH). Special thanks are due to Director, ICAR-CIBA for providing the necessary facilities and support for the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Food and Agriculture Organization (FAO). The global status of seafood safety. Rome; 2018.
2. FAO. The state of world fisheries and aquaculture 2010. Rome: FAO; 2010.
3. FAO. The state of world fisheries and aquaculture 2012. Rome: FAO; 2012.
4. Marine Products Export Development Authority (MPEDA). Annual Report 2022-23. Ministry of Commerce and Industry. Government of India; 2023
Available:<https://mpeda.gov.in/wp-content/uploads/2020/12/mpeda-annual%20report%202022-2023-2.pdf>
5. Johnson SC, Bravo S, Nagasawa K, Kabata Z, Hwang J, Ho J, Shih CT. A review of the impact of parasitic copepods on marine aquaculture. *Zoological Studies*. 2004;43(2):229-43.
6. Shinn AP, Pratoomyot J, Bron JE, Paladini G, Brooker EE, Brooker AJ. Economic costs of protistan and metazoan parasites to global mariculture. *Parasitology*. 2015;142(1):196-270.
Doi:10.1017/S0031182014001437
7. Roberts T, Hutson D. Macrocytic insecticides. In: Roberts, T., Hutson, D. (Eds.), *Metabolic Pathways of Agrochemicals: Part 2: Insecticides and Fungicides*. The Royal Society of Chemistry, Cambridge, U.K. 1999;87-94.
8. Raja RA, Patil PK, Avunje S, Aravind RP, Alavandi SV, Vijayan KK. Biosafety, withdrawal and efficacy of anti-parasitic drug *Emamectin benzoate* in Asian Seabass (*Lates calcarifer*). *Aquaculture*. 2020;30(525):735335.
DOI: 10.1016/j.aquaculture.2020.735335
9. Raja RA, Patil PK, Avunje S, Kumaran M, Solanki HG, Jithendran KP, Alavandi SV, Vijayan KK. Efficacy of *Emamectin benzoate* in controlling natural infestations of ectoparasites in economically important fish species of India. *Aquaculture*. 2022;30(551):737940.
DOI:10.1016/j.aquaculture.2022.737940
10. Halling-Sørensen BN, Nielsen SN, Lanzky PF, Ingerslev F, Lützhøft HH, Jørgensen SE. Occurrence, fate and effects of pharmaceutical substances in the environment-A review. *Chemosphere*. 1998;36(2):357-93.
DOI: 10.1016/S0045-6535(97)00354-8
11. Boreen AL, Arnold WA, McNeill K. Photodegradation of pharmaceuticals in the aquatic environment: a review. *Aquatic Sciences*. 2003;65:320-41.
DOI:10.1007/s00027-003-0672-7
12. Burrige L, Weis JS, Cabello F, Pizarro J, Bostick K. Chemical use in salmon aquaculture: a review of current practices and possible environmental effects. *Aquaculture*. 2010;15;306(1-4):7-23.
DOI: 10.1016/j.aquaculture.2010.05.020
13. Lunestad BT, Samuelsen OB, Fjelde S, Ervik A. Photostability of eight antibacterial agents in seawater. *Aquaculture*. 1995;15;134(3-4):217-25.
DOI: 10.1016/0044-8486(95)00065-A
14. Cheng B, Van Smeden J, Deneer J, Belgers D, Foekema E, Roessink I, Matser A, Van den Brink PJ. The chronic toxicity of *Emamectin benzoate* to three marine benthic species using microcosms. *Ecotoxicology and Environmental Safety*. 2020;1;194:110452.
DOI: 10.1016/j.ecoenv.2020.110452
15. Krieger MS, Yoder RN, Gibson R. Photolytic degradation of florasulam on soil and in water. *Journal of agricultural and food chemistry*. 2000;48(8):3710-7.
DOI:10.1021/jf991290+
16. Tariq MI, Afzal S, Hussain I, Sultana N. Pesticides exposure in Pakistan: A review. *Environment international*. 2007;33(8):1107-22.
DOI: 10.1016/j.envint.2007.07.012.
17. Mushtaq M, Hafeez FY, Riazuddin S. Selective photolysis of certain pesticides in aqueous environment and their photolytic degradation rate. *J. Environ. Sci. Health B. Pesticides, Food Contaminants, and Agricultural Wastes*. 1996;31(6):1205-1214.
18. Chukwudebe AC, Wislocki PG, Villeneuve DL, Sagartz JE, Crouch LS. Environmental fate of *Emamectin benzoate*: degradation in aerobic aquatic sediment systems. *J. Agric. Food Chem*. 1997;45(3):787-795.
DOI:10.1021/jf970589u

19. Keen BA, Raczkowski H. The relation between the clay content and certain physical properties of a soil. *The Journal of Agricultural Science*. 1921;11(4):441-9.
Available:<https://doi.org/10.1017/S0021859600005827>
20. Espenson JH. Chemical kinetics and reaction mechanisms. New York: McGraw-Hill; 1995.
21. Sakrabani R, Boxall A. Inputs and fate of contaminants in the aquatic environment. *Sustainable Water: Chemical Science Priorities*, Final draft; 2007.
22. Mushtaq M, Chukwudebe AC, Wrzesinski C, Allen LR, Luffer-Atlas D, Arison BH. Photodegradation of *Emamectin benzoate* in aqueous solutions. *Journal of Agricultural and Food Chemistry*. 1998;46(3):1181-91.
DOI: 10.1021/jf970561i
23. Feely WF, Crouch LS, Arison BH, VandenHeuvel WJ, Colwell LF, Wislocki PG. Photodegradation of 4''-(epimethylamino)-4''-deoxyavermectin B1a thin films on glass. *Journal of Agricultural and Food Chemistry*. 1992. 40(4):691-6.
24. Wrzesinski C, Cruz JM, Moraes J. Photolysis of ivermectin. *J. Agric. Food Chem*. 1996;44(4): 1172-1175.
25. Crouch NR, Feely WF. Effect of irradiation on avermectins in river water. *Pestic. Sci*. 1995; 45(2):169-175.
26. Yousif E, Haddad R, Greaves J. Photolysis of imidacloprid in aquatic environments: Effect of season and light intensity. *Aqua. Sci*. 2013;75(3):329-337.
27. Sachdev SL, Hannink N, Basu A, Dotson GS. Advanced oxidation processes for wastewater treatment: state of the art. *Green Chem. Lett. Rev*. 2021;14(1):22-37.
28. Mushtaq M, Hafeez FY, Riazuddin S, Malik MN. Photolysis of ivermectin in aqueous solution. *Int. J. Pharmacol*. 1995;33(2):121-125.
29. Chukwudebe A, Wislocki P, Organ R, Sparks T, Crouch L. Environmental fate of avermectin B1a in outdoor aquatic microcosms. In *Pesticide chemistry*. Elsevier. 1992;79-84.

Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/123208>