



Effect of Different Diets on the Growth Rate of the Rotifer, *Brachionus plicatilis* under 40 ppt Salinity Stress

Ahmed M. M. Heneash¹, Khalid Abd Al-Rahman² and Mohsen Y. Omer^{1*}

¹*National Institute of Oceanography and Fisheries (NIOF), National Institute of Oceanography and Fisheries, Egypt.*

²*Al-Wafa Farm of Fish Aquaculture, National Institute of Oceanography and Fisheries (NIOF), Egypt.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJEE/2021/v16i430276

Editor(s):

(1) Dr. Edward Ching-Ruey Luo, National Chi-nan University, Taiwan.

Reviewers:

(1) Quy Moc Ong, Nong Lam University, Vietnam.

(2) Adande Richard, University of Abomey-Calavi, Benin.

(3) Ojelade, Oluwaseun C., Federal University of Agriculture Abeokuta, Nigeria.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/79923>

Received 20 October 2021

Accepted 21 December 2021

Published 23 December 2021

Original Research Article

ABSTRACT

Aquaculture is important business some fish farms use underground water that may has high salinity. The aim of this study is to evaluate the effect of eight different types of diets on enhancing the growth rate of the rotifer, *Brachionus plicatilis* lives in 40 ppt saline water in 2020. The diets used in the study are T1 (*Nannochloropsis oculata*), T2 (*Tetraselmis suecica*), T3 (*N. oculata* + *T. suecica*), T4 (*N. oculata* + yeast; *Saccharomyces cerevisiae*), T5 (*T. suecica* + yeast), T6 (*S. parkle*), T7 (*S. Parkle* + yeast) and T8 (yeast). The results revealed that the rotifer fed for *S. parkle* showed the highest growth population (818 Indv. /mL) with a highly significant difference ($p < 0.05$). On the other hand, rotifers fed the yeast (*S. cerevisiae*) showed the lowest population growth (355 Indv. /mL). The outcome of the current study indicates that *S. parkle* commercial diet is very appropriate for the stock culture of *B. plicatilis* due to its ability to increase the growth rate of the rotifer in a short time under the current conditions. *S. Parkle* feeding led to an increase in the densities of rotifers in high salinities.

*Corresponding author: E-mail: my.omer@niof.sci.eg;

Keywords: *S.parkle*; yeast; *Tetraselmis suecica*; *Nannochloropsis oculata*; zooplankton; fish food.

1. INTRODUCTION

Zooplankton is the first food for many fish crops due to its effectiveness in increasing the growth rate of fish larvae. It is also considered as an easily digestible protein for fish and shellfish [1]. Rotifers as a zooplankton is used as natural food for larvae (Ichthyoplankton) in hatcheries as they are the favorite prey of most fish larvae [2].

One of the vital difficulties faced production of fish fry in hatcheries is the lack of live food for the nutrition of early stages of fish larvae. Therefore, there is no alternative food that could replace the using of rotifers as preliminary food for larvae until now [3,4]. Moreover, the rotifer, *Brachionus plicatilis* is an essential food type especially in the early larval stages of many fish species due to its small size and its ability to propagate in captivity at high density and reproductive rate [5]. Additionally, it has the ability to stand wide ranges of salinity [6].

Furthermore, rotifers need different types of food to survive and be active in the hatcheries. Various species of microalgae have been used as food for rotifers including; *Tetraselmis* spp., *Nannochloropsis* spp., *Chaetoceros* spp., *Rhodomonas* spp. and *Isochrysis* spp. [7], (Hoff and Snell 2001; and Wikfors and Ohno, 2001). In contrast, baker's yeast, *Saccharomyces cerevisiae* is immediately available compared to the microalgae (fresh, frozen or dried) which are laborious, time-consuming and expensive. Additionally, the yeast, *S. cerevisiae* has an important role to give a high population growth rate for rotifers [8]. However, *Nannochloropsis* sp. is widely used as live food for rotifers as it provides the predicted yield on time with high density because of its ease of cultivation (Abd Rahman et al, 2018).

B. plicatilis are a dominated Rotifera that are a preferred prey for some larval fish and are used by culturists during larval fish culture [9] and El-Damhogy et al., 2016. Threlkeld [10], Krebs [11], Sarma [12], Suantika et al. [13], Dhert et al. [7], Sarma et al. [14], Omar et al. [15], El-Kassas et al. [1], Heneash et al. [8] and Sharma et al. [16] reported on the culture of rotifers (*B. plicatilis*).

In Egypt, El-Wafa fish farm is located in the Ismailia governorate next to the Suez Canal and

the groundwater well is the source of aquaculture water, the well's salinity is 40 ppt in 2020. The farm produces *Dicentrarchus labrax* and *Sparus aurata*. Consequently, during the feeding of the grown fish with natural food (zooplankton), species of zooplankton that tolerate salinities above normal sea water (35-37 ppt). Hence, the present work aims to study the effects of 40 ppt salinity on the rotifer (*B. plicatilis*) growth on different feeding diets and the effect of an artificial food type and initial stocking density on the population growth of rotifers to induce high productivity of *B. plicatilis* to be used as fish food.

2. MATERIALS AND METHODS

The experiment was conducted in El-Wafa fish farm, Ismailia, Egypt. *Brachionus plicatilis* specimens were obtained from the marine hatchery of the National Institute of Oceanography and Fisheries (NIOF), Alexandria, Egypt. The rotifers were divided into eight groups each one has three replicates. Then, they were stocked in twenty-four 9- liters plastic jars supplied with continuous aeration. Water parameters were measured daily including water temperature (digital thermometer), water acidity using Jenway pH meter, (Model 350) and salinity using digital salinometer. Two marine green microalgae were tested; *Nannochloropsis oculata* and *Tetraselmis suecica*. Ecch algal species were then combined with the baker's yeast (*Saccharomyces cerevisiae*) and Selco S. parkle; which is yeast-based ready commercial diet for rotifers manufactured by INVE.

2.1 Dietary Treatments

Eight diets are (T1-T8), where T1 (*N. oculata*) was considered as a control, T2 (*T. suecica*), T3 (*N. oculata* + *T. suecica*), T4 (*N. oculata* + yeast; *S. cerevisiae*), T5 (*T. suecica* + yeast), T6 (S.parkle), T7 (S.parkle + yeast) and T8 (yeast).

2.2 Experimental Procedure

The experiment started with two liters initial volume containing 150 indiv./L initial rotifer density for each replicate in each treatment. They were supplied with the indicated diet (Table 1) twice daily at 09.00 AM and 14.00 PM in 1000 mL saline water (underground water) for 8 days.

Table 1. The eight diets of feeding diets and their feeding rates/jar/day

Treatment	Feeding diets	Initial volume of water/Jar 1 st day	Feeding rate/Jar/day
T1	<i>Nannochloropsis oculata</i>		1000 mL
T2	<i>Tetraselmis suecica</i>		1000 mL
T3	<i>N. oculata</i> + <i>T. suecica</i>		1000 mL
T4	<i>N. oculata</i> + Yeast (<i>Saccharomyces cerevisiae</i>)	1000 mL*	1000 mL+ 0.1 gm/l
T5	<i>T. suecica</i> + Yeast (<i>S. cerevisiae</i>)		1000 mL+ 0.1 gm/l
T6	S.parkle		0.1 gm/l * 50 mL
T7	S.parkle + Yeast (<i>S. cerevisiae</i>)		0.1 gm/l * 50 mL
T8	Yeast (<i>S. cerevisiae</i>)		0.1 gm/l * 50 mL

* Each one mL contains 150 indiv. /l

2.3 Sampling and Counting of *B. plicatilis*

Each zooplankton density in each treatment zooplankton was determined using a binocular microscope (40X). The density of rotifer organisms was calculated as their total number per liter. The concentrated rotifers were subsampled using a Stempel pipette [8,17,18,1].

2.4 Population (R)

Along the eight experimental days, the population of rotifer was calculated as the increase in number every 24 hours by counting the individuals (three times) in a 1-mL glass pipette. These concentrated rotifers were sampled with a Pasteur pipette and were counted under the dissecting microscope (Braley, 1994).

2.5 Population Growth Rate (R')

The population growth rate of rotifers (R') was calculated according to Okauchi and Fukusho [19] using the following equation:

$$R' = \frac{(N_t - N_0)}{t}$$

where R' = the instantaneous growth rate; N₀ = the initial number of rotifers; N_t = the final number of rotifers after t days; t = culture days.

2.6 Statistical Analysis

Data were analyzed using analysis of variance test (One-way ANOVA) to determine the significant differences among rotifer populations between treatments. Also, Duncan's multiple range test was used to provide more information of ranges of all the pairs of treatments using IBM SPSS Statistics (v22).

3. RESULTS

3.1 Water Quality

During the experiment period, the pH value was 8.0 ± 0.2. Water temperature and salinity were 28.0 ± 2.0°C, 40 ± 1.0 ppt, respectively.

3.2 Effect of Different Diets on the Growth Rates of *B. plicatilis* During the Experiment Period

During the eight experiment days, T6 (S. parkle) was the most effective diet which gave the maximum average density of *B. plicatilis*, (818 Indv. /mL). On the other hand, T8 was the least effective diet which contributed the minimum average density represented only by (355 Indv. /mL). Fig. 1 shows the impact of the eight experimental diets studied during the experiment period, indicating that the highest average value listed in the ninth day was (935 Indv./mL), while the lowest average value recorded in the first day was (265 Indv./mL). Besides, the aforementioned figure revealed that T6 was the most effective diet which gave a distinct increase in the growth rate of *B. plicatilis*.

Furthermore, Table (2) revealed that there were highly significant differences among the eight experimental diets where the significance level of (F = 2.665, p = 0.019) is less than the threshold value of (p < 0.05). In conclusion, not all the group means were equal which revealed that the effects of treatments were not similar in the present experiment.

This leads to the fact that at least one of the groups tested differs from the other. In addition, Duncan's multiple range test was used to determine the critical values for comparisons

among means. The test revealed that the treatment pairs T1 and T3, T1 and T4, T1 and T5, T1 and T6, T1 and T7, T1 and T8, T2 and T5, T2 and T6, T2 and T7, T2 and T8, T3 and T6, T3 and T7, T3 and T8, T4 and T7, T4 and T8, T5 and T7, T5 and T8 had mean differences with significant value ($p < 0.05$). The rest pairs of treatments showed a significance level greater

than (0.05) which means there were no statistically significant differences between them.

The pairwise comparison among treatments showed a highly significant difference in T6 (*S.parkle*) and T8 (*S. cerevisiae*) where p -value = 0.0015 (Fig. 2).




Fig. 1. Effect of the eight experimental diets during the experiment period

Table 2. Results of the one-way ANOVA test for the population density of rotifer for the eight experimental diets

Sources of Variation	Sum of Squares	Df	Mean Square	F	p-value
Between Groups	0.057	7	0.008	2.665	0.019
Within Groups	0.170	56	0.003		
Total	0.227	63			




Fig. 2. Boxplot provides a pairwise comparison between the experimental treatments (Duncan's multiple range test), the number of asterisks depends upon the p -value of particular pair of samples; * for $p < 0.05$ and ** for $p < 0.01$




Fig. 3. The gravid females of *B. plicatilis* (Indv. /mL) during the experiment period




Fig. 4. Schematic representation of changes in population density and egg ratio of *B. plicatilis* in relationship to time in the mean diet treatments

3.3 Gravid Females of *B. plicatilis*

The gravid females of *B. plicatilis* (Indv. /mL) during the Experiment period was shown in Fig. 3. Treatment 4 displayed the greatest average number of gravid females, (144 Indv. /mL) which began with (96 Indv. /mL) on the first day of the experiment. In contrast, the lowest average number of the gravid females of the studied rotifers was for T7 (47 Indv. /mL) which started by (43 Indv. /mL). for T6, the average number of gravid females was (57.1875 indiv. /mL) which

was (46 indiv. /mL) at the first day of the experiment.

3.4 Population Density and Egg Ratio of *B. plicatilis*

Fig. 4 illustrated the relationship between the population density of the rotifer *B. plicatilis* and the egg ratio in the present study. The population density of the studied rotifers increased exponentially with time for all the treatments and reached a noticeable level after eight days (end

of the experiment). The egg ratio decreased gradually over time during the different growth levels.

4. DISCUSSION

Rotifers are cosmopolitan organisms as they are osmoconformers and can be adapted to salinities ranging from 1‰ to 97‰, however, the changes in salinity may cause shock for rotifers affects their production rate [20,21]. Increased salinity may shorten the lifetime and decrease the fecundity of female and growth rate in some rotifer species [22]. Where Oltra and Todoli [23] found negative effect for decreasing salinity on growth rate of the marine rotifer, *Synchaeta cecilia valentine*. Numerous studies established the effect of salinity on growth rate of rotifer, *B. plicatilis* [24]; where it can tolerate wide range of salinities [25] but it has the highest reproductive rate at salinities ranging from 10- 20‰ [26]; They added that the ability of *B. plicatilis* to tolerate wide salinity range is due to its colony origin, where rotifers taken from the coastal marshes stand high salinities more than those taken from brackish water lagoons.

In the present study, the statistical analysis revealed that the eight diets showed different effects through the eight days of the experiment. The commercial diet T6 (S.parkle) Caused gradual increase in rotifer density from the third; where Lind [27] and Eryalcin, [28] recorded the highest growth rate mean for Sparkle comparing with the other rotifer diets used in his study. Furthermore, Kostopoulou et al. [29] found positive effect on the growth rate of the rotifer *B. plicatilis* with the dilution rate of S.parkle .

In contrast to the present and aforementioned findings, Qi et al. [30] found an increased growth rate of rotifer cultures fed on *Nannochloropsis* algae than S.Parkle. More importantly, Reitan and Olsen [31] found that S.parkle diet had promoted more bacteria than the algal diets; they found that yeast diet did not support growth rate of rotifers where the Yeast *S. cerevisiae* was the less effective diet in the current study; this was confirmed by the results of Planas and Estevez [32]. In contrast, [33] reported that yeast is more effective than *Chlorella* powder as combined food with fresh *Chlorella*. Sarma et al. [34] stated that the yeast was not appropriate for the rotifer cultivation but the mixture of yeast and fresh *Chlorella* could be more effective on the culture of the rotifer *B. calyciflorus* and better than the other mixtures [35,36]. Heneash et al.,

[37], reported that the best diet among the seven treatments was the yeast treatment. Attallah [38] and Suchar and Chigbu [39] observed an increase in population growth when rotifers fed higher concentration of *Nannochloropsis oculata* (1000 x 10³ Cell /ml) and (700 x 10³ Cell /ml) respectively after 7 days the population growth; while the maximum mean reproductive potential was at (100 x 10³ Cell /ml) (200 x 10³ Cell /ml).

The egg ratio was determined in the present study in order to estimate the growth rate of *B. plicatilis*. Edmondson [40] and Paloheimo [41] reported that the egg ratio was used to evaluate the fecundity and the mortality rates for deriving the population growth rates. Zaky et al., [42] stated that rotifer (*Brachionus plicatilis*) feeding on *Arthrospira platensis* was the highest significantly density of the rotifer population females carrying eggs. Ohman et al. [43] and Razlutskiy [44] had introduced a very useful bases for constant improvement in the use of the egg ratio method for zooplankton. There is usually an inverse, linear or curvilinear relationship between the density of population and the egg ratio. Duncan [45], Sarma and Rao [46] recorded that the planktonic rotifer populations continue to produce between 3 and 5 eggs per female per day at low population densities when there are plenty of resources. S. Parkle feeding led to an increase in the densities of rotifers in high salinities.

5. CONCLUSION

The present study concluded that the commercial diet S.parkle was the most effective diet among the experimented diets in the current study, because it showed the highest growth rate compared with the other rotifer diets. The study recommends feeding rotifers to S.parkle because of its positive effect on feeding rotifers in high salinities.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. El-Kassas HY, Heneash AMM, Hussein NR. Cultivation of *Arthrospira (Spirulina) platensis* using confectionary wastes for aquaculture feeding. Journal of Genetic Engineering and Biotechnology. 2015;(ISSN1687-157X)13:145–155.

2. Pfejffer TJ, Ludwig GM. Small-Scale System for the Mass Production of Rotifers Using Algal Paste. North American Journal of Aquaculture. Anrerican Fisheries Sociel). 2007;69:239-243.
3. Hagiwara A, Gallardo WG, Assavaaree M, Kotani T, de Araujo AB. Live food production in Japan: recent progress and future aspects. Aquaculture 2001;200(2-1):117-127. 117–127. 00696-2.
4. Yoshimatsu, T, Hossain MA. Recent advances in high-density rotifer culture in Japan. Aquaculture International. 2014;22(5):1587–1603. DOI:10499-014-9767-5.
5. Alam MJ, Shah MMR. Growth and reproductive performance of locally isolated brackish water rotifer (*Brachionus plicatilis*) feeding different micro algae. Bangladesh, Fish. Res. 2004;8(2):127-133.
6. Rahman ARA, Cob ZC, Jamari Z, Mohamed AM, Toda T, Ross OH. The Effects of Microalgae as Live Food for *Brachionus plicatilis* (Rotifer) in Intensive Culture System. Tropical Life Sciences Research. 2018;29(1):127–138.
7. Dhert P, Rombaut G, Suantika G, Sorgeloos P. Advancement of rotifer culture and manipulation techniques in Europe. Aquaculture. 2001;200:129–146.
8. Heneash AMM, Ashour M, Matar M. Effect of un-live microalgal diet, *Nannochloropsis oculata* and *Arthrospira (Spirulina) platensis*, comparing to yeast, on Population and Growth rate of *Brachionus plicatilis*. Mediterranean Aquaculture Journal (ISSN: 1687- 7617) 2015;7: 63 – 70.
9. Pfejffer TJ, Ludwig GM. Small-Scale System for the Mass Production of Rotifers Using Algal Paste. North American journal of Aquaculture. 2007;69:239- 243.
10. Threlkeld ST. Empty loricas and the dynamics of *Kellicottia longispina* in a subalpine, oligotrophic lake. Hydrobiologia.1983;104:367–372.
11. Krebs CJ. Ecology: The Experimental Analysis of Distribution and Abundance (3rd edn). Harper and Row, New York. 1985;800.
12. Sarma SSS. Some relationships between size structure and fertility of rotifer populations. In Kaul, B. L. (ed.), Advances in Fish and Wildlife Ecology and Biology. vol. 1. Daya Publishing House, Tri Nagar, Delhi, India.1996;37–50.
13. Suantika G, Dhert P, Nurhudah M, orgeloos PS. High-density production of the rotifer *Brachionus plicatilis* in a recirculation system: consideration of water quality, zootechnical, and nutritional aspects. Aquacultural Engineering. 2000;21:201-214.
14. Sarma SSS, Gulati RD, Nandini S. Factors affecting egg-ratio in planktonic rotifers Hydrobiologia. 2005;546:361–373. DOI 10.1007/s10750-005-4247-6
15. Omar EA, Essa MA, Srouf TM, Mater MM. Biotic and Abiotic factors Affecting Productive Performance of Rotifers (*Brachionus plicatilis*) population produced under 3 days Batch culture system J. Adv. Agric. res. 2014;19(4):800-812.
16. Sharma JG, Masuda R, Tanaka M, Chakrabarti R. The Continuous Culture of Rotifer *Brachionus plicatilis* with Seawater. Madridge J Aquac Res Dev. 2018;2(1): 40-42.
17. El-Damhoghy KA, Heneash AMM, Nasef A, Khater MM. Diversity and distribution of *Brachionus* community (Rotifera: Brachionidae) at lake Maryout, Alexandria, Egypt. International Journal of Fisheries and Aquatic Studies. 2016;4(5):500-506.
18. Aboul Ezz, Sawsan M, Heneash, Ahmed MM, Gharib, Samiha M. Variability of spatial and temporal distribution of zooplankton communities at Matrouh beaches, south-eastern Mediterranean Sea, Egypt. Egyptian Journal of Aquatic Research. 2014;40(3):283–290.
19. Okauchi M, Fukusho K. Environmental conditions and medium required for mass culture of a minute algae, *Tetraselmis tetraethele* (Prasiniphyceae). Bulletin of the National Resource Institute of Aquaculture. 1984;5:1-11.
20. Epp RW, Winston PW. Osmotic Regulation in the Brackish-Water Rotifer *Brachionus Plicatilis* (Muller). Journal of Experimental Biology. 1977;68:151-156.
21. Walker KF. A synopsis of ecological information on the saline lake rotifer *Brachionus plicatilis* Müller 1786. Hydrobiologia 1981;81:159–167.
22. Serra M, Carmona MJ, Miracle MR. Survival analysis of three clones of *Brachionus plicatilis* (Rotifera). Hydrobiologia. 1994;227:97–105.
23. Oltra R, Todolí R. Effects of temperature, salinity and food level on the life history traits of the marine rotifer *Synchaeta cecilia Valentina*, n.

- subsp. J. Plankton Res. 1997;19: 693–702.
24. Bosque T, Hernández R, Pérez R, Oltra, Todolí R. Effects of salinity, temperature and food level on the demographic characteristics of the seawater rotifer *Synchaeta littoralis* Rousselet. Journal of Experimental Marine Biology and Ecology. 2001;258:55–64.
 25. Bayly IAE. Salinity tolerance and osmotic behavior of animals in athalassic saline and marine waters. Annu. Rev. Ecol. Syst. 1972;3:233–268.
 26. Miracle, M.R., Serra, M. Salinity and temperature influence in rotifer life history characteristics. Hydrobiologia. 1989;186r187:81–102.
 27. Lind, Ryan D. Effects of Selected Commercial Diets and Yeast Substitution on the Growth and Associated Microbiota of Rotifer (*Brachionus plicatilis*); 2014.
 28. Eryalcin, Kamil Mert. Effects of Different Commercial Feeds and Enrichments on Biochemical Composition and Fatty Acid Profile of Rotifer (*Brachionus Plicatilis*, Müller, 1786 and *Artemia Franciscana*. Turkish Journal of Fisheries and Aquatic Sciences. 2018;18:81-90.
 29. Kostopoulou, Venetia, Vasilakis, Manolis and Divanach, Pascal. Semi-continuous mass culture of rotifers (*Brachionus plicatilis*) using an automatic feeder. Aquaculture Research. 2012;43:91–98.
 30. Qi Z, Dierckens K, Defoirdt T, Sorgeloos P, Boon N, Bao Z, Bossier P. Effects of feeding regime and probionts on the diverting microbial communities in rotifer *Brachionus* culture. Aquaculture International. 2009;17(4):303- 315.
 31. Reitan K, Olsen Y. Comparison of rotifer culture quality with yeast plus oil and algal-based cultivation diets. Aquaculture International. 1994;2(4):225–238.
 32. Planas M, Estevez A. Effects of diet on population development of the rotifer *Brachionus plicatilis* in culture. *Helgolander Meeresunters*. 1989;43:171-181.
 33. Khatun B, Rahman R, Rahman MS. Evaluation of Yeast *Saccharomyces cerevisiae* and Algae *Chlorella vulgaris* as Diet for Rotifer *Brachionus calyciflorus*. A Scientific Journal of Krishi Foundation, The Agriculturists. 2104;12(1): 1-9.
 34. Sarma SSS, Jurado PSL, Nandini S. Effect of three food types on the population growth of *Brachionus calyciflorus* and *Brachionus patulus* (Rotifera: Brachionidae). Revista de Biología Tropical. 2001;49(1):77-84.
 35. Ashraf M, Ullah S, Rashid T, Ayub M, Bhatti EM, Naqvi SA, Javaid M. Optimization of Indoor Production of Fresh Water Rotifer, *Brachionus calyciflorus*, b: Feeding Studies. Pakistan Journal of Nutrition. 2010;9(6): 582-588.
 36. Pena-Aguado F, Nandini S, Sarma SSS. Differences in population growth of rotifers and Cladoceras raised on algal diets supplemented with yeast. Limnologica-Ecology and Management of inland Waters. 2005;35(4): 298-303.
 37. Heneash AMM, Toutou MMM, Ahmed HO, Khafage AR. The effect of using different feeding types on the growth performance and egg production rates of Rotifers (*Brachionus plicatilis*) in marine waters. Mediterranean Aquaculture Journal. 2019;(11) 1-10.
 38. Attallah MA. Study on potential of culturing rotifer (*Brachionus plicatilis*) in an intensive closed system, MSc. Al-Azhar University. 2016:176.
 39. Suchar VA, Chigbu P. The effects of algae species and densities on the population growth of the marine rotifer, *Colurella dicentra*. Journal of Experimental Marine Biology and Ecology. 2006;337(1): 96-102.
 40. Edmondson WT. Reproductive rate of planktonic rotifers as related to food and temperature. Ecological Monographs. 1965;35:61–111.
 41. Paloheimo J. Calculation of instantaneous birth rate. Limnology and Oceanography 1974;19:692–694.
 42. Zaki MA, Ashour M, Heneash AMM, Mabrouk MM, Alprol AE, Khairy HM, Nour AM, Mansour AT, Hassanien HA, Gaber A, Elshobary ME. Potential Applications of Native Cyanobacterium Isolate (*Arthrospira platensis* NIOF17/003) for Biodiesel Production and Utilization of Its By product in Marine Rotifer (*Brachionus plicatilis*) Production. Sustainability. 2021;13:1769.
 43. Ohman MD, Aksnes DL, Runge JA. The interrelationship of copepod fecundity and mortality. Limnology and Oceanography. 1996;41:1470–1477.

44. Razlutskiy VI. Estimating cladoceran birth rate: use of the egg age distribution to estimate mortality of ovigerous females and eggs. *Hydrobiologia*. 2000;428: 135–144.
45. Duncan A. Factors influencing the composition, body size and turnover rate of zooplankton in Parakarama Samudra, an irrigation reservoir in Sri Lanka. *Hydrobiologia*. 1984;113:210–215.
46. Sarma SSS, Rao TR. The combined effects of food and temperature on the life history parameters of *Brachionus patulus* Muller (Rotifera). *Internationale Revue der gesamten Hydrobiologie*. 1991;76:225–239.

© 2021 Heneash et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/79923>