



Hepatotoxicity of Ethanol Extract of *Adenium obesum* Stem Bark in Wistar Rats

Samson Enejo Abalaka^{1*}, Muhammad Yakasai Fatihu¹,
Najume Doguwar Giginya Ibrahim¹ and Suleiman Folorunsho Ambali²

¹Department of Veterinary Pathology, Ahmadu Bello University, Zaria, Nigeria.

²Department of Veterinary Pharmacology and Toxicology, Ahmadu Bello University, Zaria, Nigeria.

Authors' contributions

This work was carried out in collaboration between all authors. Author SEA was responsible for the conceptualization and design of the work in addition to data collection, analyses and interpretations with the manuscript preparation. Author MYF was involved with the study's conceptualization and design and data interpretations. Author NDGI partook in the design of the study and data interpretations. Author SFA was also involved with the design of the study and data analysis with interpretations. All authors read and approved the final manuscript.

Original Research Article

Received 2nd November 2013
Accepted 1st February 2014
Published 20th March 2014

ABSTRACT

Aims: *Adenium obesum* is a known medicinal plant thereby creating the need for the evaluation of its toxicity and histopathological effects on the liver of female Wistar rats orally administered ethanol extract of the plant's stem bark.

Place and Duration of Study: Department of Veterinary Pathology, Ahmadu Bello University, Zaria, Nigeria, between January 2011 and January 2012.

Methodology: Ethanol extraction of *A. obesum* stem bark was performed prior to screening it for its phytochemical constituents. Female rats per group were orally administered by gavage pre-defined doses (300mgkg⁻¹, 2000mgkg⁻¹ and 5000 mgkg⁻¹) of the extract separately in a stepwise procedure and observed for signs of toxicity. Control rats were administered distilled water placebo.

Results: The extract contained some alkaloids, saponins, tannins, flavonoids, glycosides, steroids and triterpens with no anthraquinones. Exposed rats did not show signs of toxicity and neither was there any mortality. Changes in aspartate aminotransferase, alanine

*Corresponding author: Email: seabalaka@yahoo.co.uk;

aminotransferase and alkaline phosphatase activities were non-significant ($p>0.05$). Congestion and fatty degenerative changes were seen in the liver of the exposed rats, which were not significantly ($p>0.05$) different in exposed rats compared to the control.

Conclusion: *Adenium obesum* did not cause major hepatic damage in the exposed rats and therefore, it is a safe oral medicinal plant within the extract dose and exposure period used in the study.

Keywords: *Adenium obesum*; wistar rats; toxicity; biochemical parameters; histopathology.

1. INTRODUCTION

Medicinal plants are used to treat variety of ailments worldwide [1,2]. This is because medicinal herbs are integral parts of alternative therapy [3]. The World Health Organization (WHO) estimated that about 80 % of the world population presently uses herbal medicine for some aspects of their primary health care needs while plant products also play important roles in the health care system of the remaining 20%, who mainly reside in developed countries [4]. No wonder herbal medicine has attracted public attention over the past 20 years in regions where this type of medicine is easily accessible [5]. However, prolonged use of these plants is associated with toxic effects [6,7], especially as most are used indiscriminately without adequate information on their safety or toxicity risk [8]. This calls for the continuous evaluation of their toxicity in attempts to elucidate on possible risks associated with the practice.

Adenium obesum is a deciduous pachycaul shrub with half buried and distinctively swollen base along with twisted branches that bears sparse leaves, which are shed prior to the appearance of its characteristic pink “showy” flowers [9,10]. Although the plant grows mostly within the Sahel to Sudanese savannahs in Africa and also in Arabia [11,12], it is equally found worldwide where it’s cultivated for ornamental purposes [13]. The bark of the plant is chewed as an abortifacient to produce miscarriages or induce abortions [14,12] even as its latex is used to treat decaying teeth, boils and septic wounds [13,15]. Similarly, the latex and bark of the plant is used to treat bone dislocation, rheumatism, sprains, paralysis, swellings and wounds [16]. There is therefore, a need to investigate the toxicity of this medicinal plant in order to optimize its dosage, especially as pharmacology is simply toxicology at a lower dose [17] and vice versa. The fact that herbal toxicity represents a serious human health threat further makes the study very imperative [18]. Therefore, the study evaluates the toxicity and histopathological implications of the oral administration of ethanol extract of *A. obesum* stem bark on the liver of exposed Wistar rats as animal models for predicting possible effects in humans.

2. MATERIALS AND METHODS

2.1 Plant Extraction

The stems of *Adenium obesum* were gathered from the open fields of Rurum town, Rano Local Government Area, Kano State, Nigeria between the months of January – April, 2011. These were authenticated with Voucher No. 1386 at the Herbarium, Department of Biological Sciences, Ahmadu Bello University (A. B. U.), Zaria, Nigeria by Mallam Musa Mohammed. The barks were removed from the stems, sun-dried and pounded into powder before soaking 3.95 kg of it in 21 L of ethanol (96.0% vol. Sigma-Aldrich® Inc., St. Louis, MO

63178, USA) over a 72-hour period. The method of Abu-Dahab and Afifi [19] was used to concentrate the filtrate to dryness in an evaporation dish at room temperature until constant weights were obtained. The extractive yield (% w/w) of the process was calculated as described by Zhang et al. [20]. Preliminary screening of the extract for its phytochemical constituents was performed using the methods of Trease and Evans [21] and Harborne [22].

2.2 Wistar Rat Toxicity Bioassay

Female rats (169 - 189g) were obtained from the Animal Unit of National Veterinary Research Institute (NVRI), Vom, Plateau State, Nigeria. The rats were acclimatized for seven days in a well ventilated room under natural photo-period (12/12-hour) while being housed in clean metal cages. Fresh drinking water was provided *ad libitum* along with NVRI pelletized rat feed (crude protein - 19.97%, crude fibre - 6.89%, Ash - 7.60%, nitrogen free extract - 59.21% and moisture - 12.98%) during the period.

The toxicity bioassay was performed as described in the OECD guideline No. 423 [23] with the administration of pre-defined doses of the extract (300 mgkg⁻¹, 2000 mgkg⁻¹ and 5000 mgkg⁻¹) by gavage separately in a stepwise procedure with the use of three female rats per step based upon the absence of morbidity and/or mortality from preceding dose. The unexposed control rats were given distilled water placebo. Exposed rats were observed for signs of toxicity during the first 30 minutes and daily thereafter throughout the 14-day observational period. The median lethal dose (LD₅₀) of the extract was established based on the OECD guideline No. 423 [23]. Similarly, changes in body weight were used as a measure of toxicity [24].

2.3 Biochemical Analyses

Two millilitres of blood were collected from the exposed rats via vene-section under light chloroform anaesthesia at the end of the 14-day post administration of the extract. These were dispensed into sample tubes without EDTA anticoagulant and centrifuged at 1006 g for 10 minutes to obtain the serum after allowing them to clot. The Reference method by International Federation of Clinical Chemistry [25] was used to determine the aspartate aminotransferase (AST) and alanine aminotransferase (ALT) activities using an autoanalyzer (Bayer Express Plus, Model 15950, Germany.). Enzymatic hydrolysis method as described by King and Armstrong [26] was used to determine the alkaline phosphatase (ALP) activity.

2.4 Histopathological Analyses

The liver of experimental rats was harvested after sacrificing them under light chloroform anaesthesia at the end of the 14-day observational period. The harvested tissues were fixed in 10% neutral buffered formalin prior to paraffin embedding, sectioning at 5 µm and haematoxylin and eosin staining [27,28]. These were examined under light microscopy for histopathological lesions.

The nature and severity of lesions in the liver of the exposed rats were noted and determined semi-quantitatively based on the adaptation of the degree of tissue changes (DTC) method by Poleksic and Mitrovic-Tutundzic [29] and Simonato et al. [30]. This involved the progressive classification of liver alterations in stages of tissue damage where the sum of the number of lesion types within each of the three stages is multiplied by the stage coefficient to give the numerical values of the DTC using the formula: $DTC = (1 \times \sum I)$

+ $(10 \times \sum II)$ + $(100 \times \sum III)$. Alterations that did not alter the normal functioning of the liver tissue were tagged Stage I alterations. Similarly, alterations that were more severe and impaired the normal functioning of the liver were tagged stage II alterations while those that were very severe and induced irreparable liver damage were tagged stage III alterations, respectively. The grading and interpretations of the results were as follows: 0 – 10 (normal liver); 11 – 20 (slightly damaged liver); 21 – 50 (moderately damaged liver); 50 - 100 (severely damaged liver); >100 (irreversibly damaged liver).

2.5 Statistical Analyses

GraphPad Prism (GraphPad Prism, version 4.0, San Diego, California, USA.) was used to analyze the data (mean \pm SEM) where a one-way analysis of variance (ANOVA) was performed for statistical significance at $p < 0.05$, including Tukey's multiple comparison test to compare the differences between the various means. Differences between the DTC in the liver of the exposed groups and the control group were compared for statistical significance ($p < 0.05$).

3. RESULTS AND DISCUSSION

3.1 Preliminary Phytochemical Screening

The extractive yield of the extract was 6.58 % w/w. Preliminary phytochemical screening of the extract revealed the presence of alkaloids, tannins, saponins, flavonoids, glycosides, steroids and triterpenes but no anthraquinones, which are of pharmacological and toxicological importance. Similar phytochemical constituents were reported from the aqueous and methanol extracts of *A. obesum* stem bark [31,32].

3.2 Toxicity Bioassay

There were no obvious changes in the skin and fur, eyes and mucous membranes of the exposed rats and neither were there changes in their behavioural patterns. Similarly, no obvious signs of tremors, salivation, diarrhoea, sleep, convulsions and coma were seen in the exposed rats, including the absence of mortality. The LD₅₀ of the extract was therefore, greater than 5000 mgkg⁻¹ or ∞ (unclassified) based on the fixed LD₅₀ cut off values [23]. The absence of obvious signs of toxicity, including mortality was indicative of the very low toxicity of the extract in the exposed rats resulting in the obtained very high LD50 value. This is in spite of the fact that the plant is a potent arrow poison [13,33]. The toxicity of the plant might therefore, be influenced by the route of administration as animals are normally exposed parenterally when the plant is used as arrow poison unlike the oral route of administration of the present study. This in addition to the fact that the toxicity of the plant is influenced by the age and parts of the plant used, genetic variation between species, climatic conditions and the soil profile of where the respective plants are found [34,35].

There were gains in body weight of experimental rats but this was significant ($p < 0.05$) only in the unexposed control rats as shown in Fig. 1. Therefore, the extract did not negatively affect the growth of exposed rats, indicative of its very low toxic nature in the exposed rats. This is because toxic chemicals or drugs adversely affect growth or weight gain in exposed animals [24].




Fig. 1. The effects of oral administration of ethanol extract of *Adenium obesum* stem bark on body weights of the exposed Wistar rats

* Significantly ($p < 0.05$) different from its control

3.3 Biochemical Analyses

Changes in some biochemical parameters of the exposed rats are as shown in Fig. 2. Although there were decreases in AST and ALP activities with increased ALT activity in the exposed rats, these changes were non-significant ($p > 0.05$) and also not concentration-dependent. The non-significant ($p > 0.05$) changes in ALT activity showed that the extract did not cause considerable damage in the liver of the exposed rats. This is because serum enzymes are cytoplasmic and are only released into circulation in cellular damage [36] where ALT activity is more hepato-specific than AST activity [37]. Similarly, the non-significant ($p > 0.05$) changes in ALP activity showed that the extract did not cause hepatobiliary problems. The toxicological importance of the decreased AST activity is unknown [38] and well less understood compared to the significance of its increased activity [39]. Mgbojikwe [31] reported similar non-significant ($p > 0.05$) changes in the AST, ALT and ALP activities of Wistar rats topically exposed to the aqueous extract of *A. obesum* stem bark.




Fig. 2. The effects of oral administration of ethanol extract of *Adenium obesum* stem bark on aspartate aminotransferase, alanine aminotransferase and alkaline phosphatase activities of the exposed Wistar rats.

3.4 Histopathological Analyses


Histopathological lesions were seen in the liver of the exposed and unexposed rats but these were comparatively to a lesser extent in the unexposed rats. The liver of the exposed rats showed congestions of the central vein along with hepatic fatty degenerations as shown in Figs. 3 to 6 with a cumulative DTC value of 3.33 ± 1.67 , indicative of the normal functioning of the liver based on the DTC grading. The incidence of histopathological lesions in the liver of the exposed and unexposed rats is as shown in Table 1. However, there were no significant ($p > 0.05$) differences between the DTC in the liver of the exposed groups compared to that of the unexposed control group.

The observed congestion and fatty degenerative changes might be due to the unique vascular, secretory, synthetic and metabolic features of the liver [40, 41]. This is because of its ability to degrade toxic compounds but can easily be overwhelmed by elevated concentrations of these compounds resulting in its structural damage [42]. The hepatic fatty degeneration is indicative of metabolic disturbance, which is a normal feature of toxic exposures [43]. These changes are usually reversible except in some extreme cases where the functional efficiency of the affected liver might be compromised [44]. Similar congestion and fatty degenerative changes were reported in the liver of Wistar rats exposed to extracts of *Sorghum bicolor* leaf sheath [45]. The fact that the DTC in the liver of exposed groups compared to the unexposed control group was non-significant ($p > 0.05$) showed that the extract did not cause considerable liver damage or hepatotoxicity in the exposed rats.

Table 1. The incidence of degree of tissue changes (DTC) in the liver of Wistar rats administered ethanol extract of *Adenium obesum* stem bark

Histopathological lesions	DTC stage	Extract dose			
		0 (Control)	300 mgkg ⁻¹	2000 mgkg ⁻¹	5000 mgkg ⁻¹
Vacuolations	I	0	0	0	+
Congestion	II	0	+	+	+

(0) – absent; (+) – rare; (++) – low incidence; (+++) – high incidence



**Fig. 3. Photomicrograph of the liver of Wistar rats administered distilled water placebo (unexposed control). Note the central vein (V) and the hepatocytes (arrows).
H & E x 397**




Fig. 4. Photomicrograph of the liver of Wistar rats administered 300 mgkg^{-1} of ethanol extract of *Adenium obesum* stem bark. Note the central vein (C) and the hepatocyte (arrow). H & E x 397




Fig. 5. Photomicrograph of the liver of Wistar rats administered 2000 mgkg^{-1} of ethanol extract of *Adenium obesum* stem bark. Note the hepatocyte (arrow) and the central vein (V) that is congestion (C). H & E x 397




Fig. 6. Photomicrograph of the liver of Wistar rats administered 5000 mgkg⁻¹ of the ethanol extract of *Adenium obesum* stem bark. Note the central vein (V) with congestion (C) and vacuolation of the hepatic cells (arrows). H & E x 397

4. CONCLUSION

The ethanol extract of *A. obesum* stem bark did not cause major liver damage in the exposed rats and therefore, might be a safe oral medicinal plant within the limitations of the study's extract dose and exposure period. However, there is a need for further investigation over repeated and prolonged exposures.

CONSENT

Not applicable.

ETHICAL APPROVAL

All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable. All experiments have been examined and approved by the appropriate ethics committee.

ACKNOWLEDGEMENTS

The authors want to appreciate the efforts of Mr. Abdulsalam Umar towards the work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Hostettmann K, Marston A, Ndjoko K, Wolfender J. The potential of African plants as a source of drug. *Curr Org Chem*. 2000;4:973-1010.
2. Ahmad I, Agil F, Owais M. *Modern phytomedicine: Turning medicinal plants into drugs*. West-Sussex. England: John Wiley and Sons. 2006;2-22. ISBN 9783527315307.
3. Karim A, Nouman M, Munir S, Sattar S. Pharmacology and phytochemistry of Pakistani herbs and herbal drugs used for treatment of diabetes. *Int J Pharmacol*. 2011;7:419-439.
4. Hoareau L, Da silva EJ. Medicinal plants: A re-emerging health aid. *Electronic J Biotechnol*. 1999;2(2):56-70.
5. Humber JM. The role of complementary and alternative medicine: Accommodating pluralism. *J Am Med Assoc*. 2002;288:1655-1656.
6. Tédonga L, Dzeufiet PDD, Dimo T, Asongalem EA, Sokeng SN, Flejou JF, et al. Acute and subchronic toxicity of *Anacardium occidentale* Linn (*Anacardiaceae*) leaves hexane extract in mice. *Afr J Trad Compl Altern Med*. 2008;4(2):140-147.
7. Ilodigwe EE, Akah PA, Nworu CS. Evaluation of the acute and subchronic toxicities of *Spathodea campanulata* P. Beauv. *Int J Appl Res Nat Prod*. 2010;3(2):17-21.
8. Ouédraogo M, Zerbo P, Konaté K, Barro N, Sawadogo LL. Effect of long-term use of *Sida rhombifolia* L. extract on haemato-biochemical parameters of experimental animals. *Br J Pharmacol Toxicol*. 2013;4(1):18-24.
9. McLaughlin J, Garofalo J. *Desert Rose (Adenium obesum)*. Miami-Dade County/University of Florida Cooperation Extension Services Fact-Sheet No. 17. 2002; Accessed 21 December 2010. Available: http://www.miami-dade.ifas.ufl.edu/pdfs/ornamental/ornamental_publications/desert-rose.PDF
10. Zorloni A. Evaluation of plants used for the control of animal ectoparasitosis in southern Ethiopia (Oromiya and Somali regions). Degree of Magister Scientiae Dissertation, South Africa: University of Pretoria; 2007.
11. Plaizier AC. A Revision of *Adenium* Roem and Schult and of *Diplorhynchus* Welw. Ex Fic. & Hiern (*Apocynaceae*). Wageningen, Netherlands: Mededelingen Land bouwhoge school, Publication No. 80-12. 1980;1-40.
12. Arbonnier M. *Trees, Shrubs and Lianas of West African Dry Zones*. MNHN: CIRAD, Margraf Publishers GMBH. 2004;161. ISBN 9782876145795.
13. Oyen LPA. *Adenium obesum* (Forssk.) Roem. & Schult. In: Schmelzer GH, Gurib-Fakim A, editors. *Plant Resources of Tropical Africa 11 (1): Medicinal Plant 1*. Wageningen, Netherlands: PROTA Foundation/Backhuys Publishers/CTA. 2008;46-49. ISBN 9789057822049/9783823615316.
14. Neuwinger HD. *African Traditional Medicine: A Dictionary of Plant Use and Applications*. Stuttgart, Germany: Medpharm Scientific. 2000;589. ISBN 9783887630867.
15. Bawden-Davies J. *The Adenium Species*; 2010. Accessed 21 December 2010. Available: http://www.ehow.com/about_6721472_adenium-species.html
16. Anonymous. *Plant Story: Desert Rose (Adenium obesum)*; 2011. Accessed 9 August 2011. Available: http://www.kew.org/science-conservation/save-seed-prospect/millenium-seed-bank/using-our-seeds/helping-communicate-orlwide/useful-plants-project/adenium-obesum/UPP-Adenium_obesum.htm.
17. Sasidharan S, Darah I, Jain K. *In vivo* and *in vitro* toxicity study of *Gracilaria changii*. *Pharm. Biol*. 2008;46:413-417.
18. Chen XW, Serag ES, Sneed KB, Zhou SF. Herbal bioactivation, molecular targets and the toxicity relevance. *Chem Biol Interact*. 2011;192(3):161-176.

19. Abu-Dahab R, Afifi F. Anti-proliferative activity of selected medical plants of Jordan against a breast adenocarcinoma cell line (MCF7). *Sci Pharm*. 2007;75:121-136.
20. Zhang S, Bi H, Liu C. Extraction of bio-active components from *Rhodiola sachalinensis* under high hydrostatic pressure. *Separ Purif Tech*. 2007;57:275-280.
21. Trease GE, Evans WC. *Pharmacognosy*. 12th ed. Philadelphia: Baillière Tindall, USA. 1983;338-628. ISBN 9780702010071.
22. Harborne JB. *Phytochemical Methods: A Guide to Modern Techniques of Plants Analysis*. Third edition. London: Chapman & Hill. 1998;279. ISBN 0412572605/0412572702.
23. Organisation for Economic Co-operation and Development (OECD). Guidelines for the Testing of Chemicals No. 423: Acute Oral Toxicity – Acute Toxic Class Method (adopted: 17th December, 2001). Paris: Organisation for Economic Co-operation and Development. 2001;1-14.
24. Teo S, Stirling D, Thomas S, Hoberman A, Kiorpes A, Khetani V. A 90-day oral gavage toxicity study of D-methylphenidate and D, L-methylphenidate in Sprague-Dawley rats. *Toxicol*. 2002;179(3):183-196.
25. Schwartz MK, De Cediell N, Curnow DH, Fraser CG, Porter CJ, Worth HG, et al. International Federation of Clinical Chemistry, Education Committee and International Union of Pure and Applied Chemistry, Division of Clinical Chemistry: Definition of the terms certification, licensure and accreditation in clinical chemistry. *J Clin Chem Clin Biochem*. 1985;23(12):899-901.
26. King EJ, Armstrong AR. A convenient method for determining serum and bile phosphatase activity. *Can Med Assoc J*. 1934;31:376-381.
27. Roberts RJ. The patho-physiology and systemic pathology of teleost. In: Roberts RJ, editor. *Fish Pathology*. London: Baillière Tindall. 1978;55-91. ISBN 9780702006746.
28. Bancroft JD, Cook HC. *Manual of histological techniques and their diagnostic application*. London: Churchill Livingstone. 1994;289-305. ISBN 9780443045349.
29. Poleksic V, Mitrovic-Tutundzic V. Fish gills as a monitor of sublethal and chronic effects of pollution. In: Mulle R, Llyod R, editors. *Sublethal and chronic effects of pollutants on freshwater fish*. Oxford: Fishing News Books. 1994;339-352. ISBN 9780852382073.
30. Simonato JD, Guedes CLB, Martinez CBR. Biochemical, physiological and histopathological changes in the neotropical fish *Prochilodus lineatus* exposed to diesel oil. *Ecotoxicol Environ Saf*. 2008;69:112-120.
31. Mgbojikwe LO. Acaricidal properties of the aqueous stem bark extract of *Adenium obesum*. PhD Dissertation. Jos University of Jos; 2000.
32. Tijjani A, Sallau MS, Sunusi I. Synergistic activity of methanol extract of *Adenium obesum* (*Apocynaceae*) stem-bark and oxytetracycline against some clinical bacterial isolates. *Bayero J Pure Appl Sci*. 2011;4(1):79-82.
33. Jones DE. *Poison arrows: North American Indian hunting and warfare*. Austin: University of Texas Press. 2007;27. ISBN 9780292714281.
34. Norton S. Toxic effects of plants. In: Klaassen CD, editor. *Casarett and Doull's toxicology: The basic science of poison*. 5th ed. New York, USA: McGraw-Hill. 1975;841-853. ISBN 9780071054768.
35. Sellappan S, Akoh CC, Krewer G. Phenolic compounds and antioxidant capacity of Georgia-grown blueberries and blackberries. *J Agric Food Chem*. 2002;50:2432-2438.
36. Sallie R, Tredger RS, Williams F. *Drugs and the liver*. Biopharma Drug Dispos. 1991;12:251-259.
37. Herfindal DR, Gourley ET. *Textbook of therapeutic drug and disease management*. 7th ed. Philadelphia, USA: Lippincott Williams and Wilkins. 2000;83. ISBN 9780781724142.

38. Ambali FS, Akanbi DO, Shittu M, Giwa A, Oladipo OO, Ayo JO. Chlorpyrifos-induced clinical, haematological and biochemical changes in Swiss albino mice – Mitigating effect by co-administration of vitamins C and E. *Life Sci J.* 2010;7(3):37-44.
39. Nanji AA. Decreased activity of commonly measured serum enzymes: Causes and clinical significance. *Am J Med Technol.* 1983;49(4):241-245.
40. Jaeschke H, Gores GJ, Cederbaum AI, Hinson JA, Pessayre D, Lemasters JJ. Mechanisms of hepatotoxicity. *Toxicol Sci.* 2002;65(2):166-176.
41. Modi H, Patel V, Patel K. Hepatoprotective activity of *Aegle marmelos* against ethanol induced hepatotoxicity in rats. *J Pharma Clin Res.* 2012;5(4):164-167.
42. Brusle J and Gonzalez G. The structure and function of fish liver. In: Munshi JSD, Dutta HM, editors. *Fish morphology.* India: Science Publishers Inc; 1996. ISBN 9054102896.
43. Wolfe JC, Wolfe MJ. A brief overview of nonneoplastic hepatic toxicity in fish. *Toxicol Pathol.* 2005;33(1):75-85.
44. Cotran RS, Kumar V, Collins T. Cellular pathology II: Adaptions, intracellular accumulations and cell aging. In: Cotran RS, Kumar V, Collins T, editors. *Robbins's pathological basis of diseases.* 6th ed. Philadelphia: WB. Saunders Company. 1999;39. ISBN 9780721673356.
45. Ogunka-Nnoka CU, Uwakwe AA, Nwachoko NC. Serum enzyme and histological studies of albino rat treated with ethanol/potash extract of *Sorghum bicolor* leaf sheath. *Indian J Drugs Dis.* 2012;1(3):74-78.

© 2014 Abalaka et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history.php?iid=464&id=14&aid=4070>